**NOVEMBER 15, 1949** 

# MODERN MEDICINE

The Journal of Diagnosis and Treatment

Papillary Adenocarcinoma of the Kidney (see page 57)

Table of contents



Dr. W. Calhoun Stirling and Col. J. E. Ash (see page 8)

Sec. 34 66, P. L. & R. U S. POSTAGE PAID PERMIT No. 29 Long Projrie, Minn.

MODERN MEDICINE 84 S. 10 St., Minneapolis 3, Minn

POSTMASTER: If undeliverable FOR ANY REASON, notify sender, stating reason on form 3547.

DNIVERSITY MICROFILMS
NO FIRST ST
NO FIRST ST
NO FIRST ST
NO FIRST ST



# Gives the cough relief your patient wants...

# Leaves the cough reflex he needs





In the average case, it's usually possible to control the patient's cough—but often it's a real problem to do it without impairing the cough reflex he needs to keep bronchioles and throat passages clear. That's where you'll find pleasant-tasting Mercodol unique!

For Mercodol contains the cough-controlling narcotic<sup>1</sup> that gives better antitussive action than codeine or heroin, yet keeps beneficial cough reflex... a superior bronchodilator<sup>2</sup> to relax plugged bronchioles... an effective expectorant<sup>3</sup> to liquefy secretions. And you'll find Mercodol notably free from nausea, constipation, retention of sputum, and cardiovascular and nervous stimulation.

# MERCODOL<sup>®</sup>

AN EXEMPT NARCOTIC

The antitussive syrup that controls cough—keeps the cough reflex



Each 30 c.c. contains:

¹Mercodinone ° 10.0 mg.

²Nethamine ® 0.1 gm.
³Sodium Citrate 1.2 gm.
°Trademark.





In tests on 2077 newborn infants . . . routine skin care with Lotion 10FA,\*





which leaves a discontinuous film, reduced incidence of miliaria to a low;



FREE! Mail coupon for 12 distribution samples!

Johnson & Johnson, Baby Products Div.
Dept. D6, New Brunswick, N. J.

Please send me 12 free distribution samples of Johnson's Baby Lotion.

Name

Street

City State

Offer limited to medical profession in U.S.A.


indicating the superiority of Lotion 10FA\* for nursery or home use.



\*Available commercially as

# JOHNSON'S BABY LOTION

Johnson-Johnson



VIM syringes are tested to withstand 20% to 40% greater pressure without leakage than government standards require. Markings are easily read, the action is smooth, and the complete syringe is annealed three times to prevent breakage in sterilization. These facts explain why VIM syringes out-perform and out-last. Specify...






hypodermic needles and syringes

MACGREGOR INSTRUMENT COMPANY, NEEDHAM 92, MASS

# Beyond the reach of human fingers



Beneficial effects may be exerted, not just locally but systemically, "beyond the reach of human fingers" in such conditions as arthritis, myositis, muscle sprains, bursitis and arthralgia. That systemic as well as local effects may be achieved by such preparations as Baume Bengué was conclusively demonstrated by the fundamental work of Moncorps, Kionka, Hanzlik, Brown and Scott.



LOCALLY-at the site of discomfort analysesic relief and a beneficial hyperemia may be readily induced.

**SYSTEMICALLY**—the salicylate absorption promoted by Baume Bengue's methyl salicylate concentration produces systemic effects to reinforce other indicated therapeutic measures.

Baume Bengué provides 19.7% methyl salicylate, 14.4% menthol in a specially prepared lanolin base.

Baume Benqué ANALGÉSIQUE

THOS. LEEMING & CO., INC., 155 E. 44th ST., NEW YORK 17

# MODERNE MEDICINE

THE JOURNAL OF DIAGNOSIS AND TREATMENT

#### EDITORIAL BOARD

James T. Case, M.D., Chicago Franklin D. Dickson, M.D., Kansas City Julius Hess, M.D., Chicago Walter B. Hoover, M.D., Boston Foster Kennedy, M.D., New York City John C. Krantz, Jr., PH.D., Baltimore A. J. Lanza, M.D., New York City Milton S. Lewis, M.D., Nashville George R. Livermore, M.D., Memphis Francis W. Lynch, M.D., St. Paul Cyril M. MacBryde, M.D., St. Louis Karl A. Meyer, M.D., Chicago Alton Ochsner, M.D., New Orleans Robert F. Patterson, M.D., Knoxville Edwin B. Plimpton, M.D., Los Angeles Fred W. Rankin, M.D., Lexington, Ky. John Alton Reed, M.D., Washington Rufus S. Reeves, M.D., Philadelphia Dalton K. Rose, M.D. St. Louis Howard A. Rusk, M.D., New York City Roger S. Siddall, M.D., Detroit James S. Simmons, M.D., Boston W. Calhoun Stirling, M.D., Washington Frank P. Strickler, M.D., Louisville Richard Torpin, M.D., Augusta, Ga. Paul M. Wood, M.D., New York City

SCIENCE WRITERS: Thomas B. Gibbons, M.D., Jorma M. Leinassar, M.D., Bernardine Lufkin, Patricia O. Read, Elizabeth Kane, Edward S. Niederkorn, David W. Morgan

TRANSLATORS: John W. Brennan, M.D., Julius Lichtenstein

# EDITORIAL CONSULTANTS

Carolyn G. Adams, M.D. E. R. Anderson, M.D. Joe W. Baird, M.D. Edward P. Burch, M.D. James B. Carey, M.D. C. D. Creevy, M.D. C. J. Ehrenberg, M.D. W. K. Haven, M.D. Charles Hymes, M.D. Miland E. Knapp, M.D. Ralph T. Knight, M.D. John F. Pohl, M.D. Leo Rigler, M.D. Wallace P. Ritchie, M.D. M. B. Sinykin, M.D. A. V. Stoesser, M.D. Arthur L. H. Street, LL.B. Marvin Sukov, M.D. Harry A. Wilmer, M.D.

#### EDITORIAL STAFF

A. E. Hedback, M.D. Editor Thomas Ziskin, M.D.

Associate Editor

Maurice B. Visscher, M.D.

Consultant in Sciences

Mark S. Parker Executive Editor

Sarah A. Davidson Managing Editor

Daisy Stilwell
Art Editor

James Niess
Editorial Board
Secretary

Margaret W. Brodt Eva Murray Editorial Assistants emotional equilibrium for the geriatric patient



Thousands of doctors know that Benzebar\* allays the mental depression, apprehension and nervousness so frequently found in the elderly patient. 'Benzebar' is S.K.F.'s logical combination of Benzedrine\* Sulfate (racemic amphetamine sulfate, S.K.F.) and phenobarbital. It provides the unique improvement of mood of 'Benzedrine' Sulfate and the calming, soothing influence of phenobarbital. These two established agents work together to restore the elderly person's zest for life and living, to quiet his nervousness and, at the same time, to keep him from overdoing.

Smith, Kline & French Laboratories, Philadelphia

Benzebar for the depressed and nervous patient




Contents for November 15 1949

VOL. 17, NO. 22

LETTER FROM THE EDITOR 14
CORRESPONDENCE. 16
WASHINGTON LETTER
QUESTIONS & ANSWERS 42
FORENSIC MEDICINE
SPECIAL ARTICLE
Papillary Adenocarcinoma of the Kidney W. Calhoun Stirling and J. E. Ash 57
MEDICINE
Potassium Deficiency J. W. Hollingsworth 61
Cracking Joints  Edward E. Brown
Intramuscular Quinidine Gluconate 62
Endocrine Treatment of Alcoholism  John W. Tintera and Harold W. Lovell 63
Rennin Estimation 64
Incidence of Hypertension 64
Radioactive Iodine in Thyroid Diagnosis D. G. Arnott, E. W. Emery, Russell Fraser, and Q. J. G. Hobson 65
Hepatic Insufficiency
GYNECOLOGY
The Menopausal Syndrome Ernest Hock



THE MEN ON THE COVER are W. Calhoun Stirling, M.D., Washington, D. C., of MODERN MEDICINE Editorial Board, and Col. J. E. Ash, M.C., U.S.A., retired, long-time director of the Army Institute of Pathology, Washington, D. C. Dr. Stirling, an international authority on genitourinary surgery, and Col. Ash, a preeminent pathologist, collaborated on the Special Article, "Papillary Adenocarcinoma of the Kidney" which starts on page 57.



Liver-stomach concentrate

Iron

vitamin B complex

... these are known raw materials for erythrocyte maturation. All are contained in Pulvules 'Lextron F.G.' (Liver-Stomach Concentrate with Ferrous Gluconate and Vitamin B Complex, Lilly). "F.G." refers to ferrous gluconate, a well-tolerated iron salt preferred by many clinicians. Pulvules 'Lextron F.G.' prescribed according to individual requirements will adequately treat any type of anemia which responds to liver or iron therapy. Available in bottles of 84 or 500.



Lilly Indianapolis 6, Indiana, U.S.A. ELI LILLY AND COMPANY



Contents

for

November 15

1949

CONTINUED

## SURGERY

Safe Induction of Pneumothorax  I. J. Selikoff, I. G. Tchertkoff, and	
E. H. Robitzek	67
Anatomy and Repair of Varicose Veins R. Stanton Sherman	68
Peptic Ulcer following Splanchnicectomy Stephen C. Mason and H. M. Pollard.	70
Physiologic Basis for Burn Treatment Carl A. Moyer	71
ORTHOPEDICS	
Traction and Suspension for Fractures Frank E. Stinchfield	74
PEDIATRICS	
Roseola Infantum William Berenberg, Charles A. Janeway, and Stanley Wright	79
Bronchiectasis in Childhood  C. Elaine Field	80
Tonsils, Adenoids, and Allergy Norman W. Clein	81
Development of Vision in Childhood  Arnold Gesell	82
PSYCHIATRY	
Histamine Therapy with Electric Shock A. M. Sackler, M. D. Sackler, and R. R. Sackler	83
VENEREOLOGY	
Office Treatment of Syphilis  Theodore Rosenthal	84
DERMATOLOGY	
Sarcoidosis	
Disseminated Neurodermatitis	85
OPHTHALMOLOGY	
The Eye in General Practice A. J. Ballantyne	86
Surgery for Primary Glaucoma Conrad Berens, L. Benjamin Sheppard, Arthur B. Duel, Jr., and Louis J. Girard	88

# multiple action without reaction

Bacteriostatic, demulcent and detergent in its positive actions, ARGYROL constantly demonstrates its advantages for effective control of infection and restoration to normal function.

Additionally, its use does not handicap the restoration process by compensatory congestion, the experience so often suffered with many vasoconstrictors.

#### The ARGYROL Technique

- The nasal meatus...by 20 per cent ARGYROL instillations through the nasolacrimal duct.
- 2. The nasal passages . . . with 10 per cent ARGYROL solution in drops.
- The nasal cavities . . . with 10 per cent ARGYROL by nasal tamponage.

## Its Three-Fold Effect

- 1. Decongests without irritation to the membrane and without ciliary injury.
- 2. Definitely bacteriostatic, yet non-toxic to tissue.
- 3. Stimulates secretion and cleanses, thereby enhancing Nature's own first line of defense.

- the medication of

choice in treating para-nasal infection. SPECIFY THE ORIGINAL ARGYROL PACKAGE

> Made only by the A. C. BARNES COMPANY NEW BRUNSWICK, N. J.

ARGYROL is a registered trademark, the property of A. C. Barnes Company



Contents for

November 15

1949

CONTINUED

UROLOGY	
Sterile Urinary Drainage  Weston T. Buddington and  Roger C. Graves	q
Cause and Therapy of Pelvic Myalgia Norborne B. Powell	0
NEUROLOGY	
Bell's Palsy  Jerome A. Hilger	
Cerebral Apoplexy Harry M. Zimmerman	04
RADIOLOGY	
Sigmoid Volvulus	3
Bronchial Stones95	3
MEDICAL FORUM	
Gastric Suction after Cesarean Delivery 94	1
Ménière's Syndrome94	1
Correction of Hypospadias98	
Retropubic Prostatectomy100	,
Heparin Therapy of Coronary Thrombosis 102	
Liver Function of Chronic Alcoholics104	ı
Giant Cell Tumor of Bone107	
Acute Rheumatic Disease in Children 108	
DIAGNOSTIX	
MEDICAL NEWS	
Interpreting Medicine for the Layman Steven M. Spencer114	
SHORT REPORTS128	
CURRENT BOOKS & PAMPHLETS152	
PATIENTS I HAVE MET	

MODERN MEDICINE, The Journal of Medical Progress, is published twice monthly on the first and fifteenth of each month at Minneapolis, Minn. Subscription rate: \$5.00 a year, 25c a copy. Business Manager: M. E. Herz. Address editorial correspondence to 84 South 10th Street, Minneapolis 3, Minn. Telephone: Bridgeport 1291. Advertising Representatives: New York 17: George Doyle, Bernard A. Smiler, Lee Klemmer, 1 East 42nd Street, Suite 801, Corn Exchange Bank Bldg. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Murray Hill 2-8717. Telephone: Murray Hill 2-8717. Teleph phone: Central 6-4619. San Francisco 4: Duncan A. Scott & Co., Mills Bldg. Telephone: Garfield 1-7950. Los Angeles 5: Duncan A. Scott & Co., 2978 Wilshire Blvd. Telephone: Dunkirk 8-4151.



So palatable and so readily digestible is LIVIBRON, nutrient hematinic containing ferrous iron, liver concentrate, and vitamin supplements, that it is tolerated readily by even the most dyspeptic of patients. These are qualities which so eminently adapt it for use in senescence, during pregnancy, and through convalescence following surgery or debilitating illness.

Nutrient tonic and hematinic effects of LIVIBRON specifically offset post-illness asthenia. LIVIBRON may be used advantageously also to meet added vitamin and hematinic requirements of pregnancy and as a general supportive measure in the aged. The pleasant flavor of LIVIBRON assures ready acceptance by children too.

Each fluid ounce of LIVIBRON contains: liver concentrate equivalent to 10 Gm. fresh liver; thiamine 5 mg.; riboflavin 2 mg.; ferrous sulfate 12 gr.; manganese citrate ¼ gr.

LIVIBRON: Supplied in 8 ounce and 1 gallon bottles.

PARKE, DAVIS & COMPANY



## LETTER FROM THE EDITOR

Dear Reader:

The other day we received a note from a doctor who wrote:

I would like to know how it is that I have been receiving copies of your publication. I do not recall having subscribed for it, but would surely do so if you will advise me of my renewal date. Please let me know if I owe you anything.

Perhaps you, too, have wondered about the same thing.

The answer is simple.

We endeavor to send Modern Medicine to every practicing physician in the United States. Our function is to perform a selective reading service for all physicians and to report the developments in medicine that are of interest to a much wider group of practitioners than the regional and specialty journals can hope to reach. To make this service effective and a real contribution to the practice of medicine it should reach as many physicians as possible. Most medical journals are distributed through subscription. This is perfectly proper, for they appeal to a limited group. However, circulation by subscription is expensive to maintain and must always fall short of complete coverage. At the start of publication seventeen years ago, Modern Medicine decided to forego revenue from subscriptions in the interest of reaching the greatest number of doctors. The record of the years testifies to the wisdom of this decision.

But the journal is free only in the sense that the reader does not have to pay for it. Editorial talent, slick paper, engravings, printing, and postage cost money. This money comes from the sale of advertising. Most journals derive their income from the sale of advertising and subscriptions. Modern Medicine's comes from the advertisers alone. Not from a few, but from many. Thus the advertisers, hundreds of them, foot the bill.

All that you are asked to do is to read MODERN MEDICINE. If you do that, all of us will feel amply repaid.

Enticement..

to



# SUSTAINED PENICILLIN LEVELS

when you prescribe penicillin

# CONFETS

[FLAVORED TABLETS BUFFERED PENICILLIN SCHENLEY]

for finicky young patients

Each tasty CONFET supplies 50,000 units of crystalline penicillin G potassium buffered with calcium carbonate. These flavored, gaily colored tablets look and taste like candy...make maintenance therapy as welcome as a reward.

SCHENLEY LABORATORIES, INC.

Executive Offices: 350 Fifth Avenue, New York 1, N. Y.

SUPPLIED: Glass tubes containing 12 tablets, 50,000 units each, stable at room temperature; no refrigeration required.









# Correspondence

Communications from the readers of Modern Medicine are always welcome. Address communications to The Editors of Modern Medicine, 84 South 10th St., Minneapolis 3, Minn.

#### Sunburn Protection

TO THE EDITORS: I find your magazine very valuable and a timesaver. Recently I've been going through some old copies and in the Oct. 1, 1948 number in the "Short Reports" I find an item about two compounds advocated by Drs. W. D. Kumler and T. C. Daniels for protection against sunburn (p. 92). I believe the reference was inadvertently omitted. Or if the work hadn't been published, do you know whether it has since appeared in print in more detail than you gave? I want to find out whether the products are on the market and if so under what name.

On a few other occasions the source of your material has not been stated. This is my only adverse criticism.

STELLA SIKKEMA, M.D.

Boulder, Colo.

¶The compounds used by Drs. Kumler and Daniels (ethyl-p-diethylaminobenzoate and methyl-p-dimethylaminobenzoate) are not yet commercially available. As to Dr. Sikkema's criticism, sources are always given if the work has been previously published.—Ed.

#### Read by Nurse, Too

TO THE EDITORS: May I tell you how much my nurse and I enjoy your magazine.

BEN EINHORN, M.D.

Los Angeles

#### Insulin and Obese Diabetics

TO THE EDITORS: In your Medical Forum section recently there appeared a comment from Dr. Arthur R. Colwell on the insulins and diabetes (July 1, 1949, p. 73). This article referred to the Symposium on Diabetes which appeared in your June 1 issue and which, incidentally, was quite complete.

Dr. Colwell lists obese individuals under the caption of those who should have protamine zinc insulin. With this listing I must take issue. This subject has been discussed with Dr. Colwell and I know that he does not intend to convey the meaning which is conveyed. It is difficult to recall any severe obese diabetics in my twenty-five years of seeing patients.

I am sure he does not refer to the older individuals because he has another group so designated. Furthermore, under the caption of globin insulin and protamine insulin mixtures, he lists young individuals. One is more or less forced to the conclusion, therefore, that obese diabetics should be given protamine zinc insulin.

While I am sure Dr. Colwell will insist upon weight reduction, I think more emphasis should be placed upon the state of obesity, particularly in the older or middle-aged patients who



# LITTLE OR NO ITCHING! LITTLE OR NO IRRITATION!

In 1948, this basically new product won nation-wide acclaim . . . Physicians, nurses and hospital authorities welcomed SEAMLESS PRO-CAP ADHESIVE PLASTER as a scientific advance that has long been needed . . . Thousands of tests have proven that PRO-CAP is the answer to those annoying and trouble-some problems that have long been an obstacle to physicians and nurses . . . A notable success because it fills a widespread and serious need . . . Sold only through leading surgical and hospital supply dealers . . . Write now for illustrated brochure and reprints of medical reports.

FINEST QUALITY SINCE 1877



# DESTRUCTION OF RECTAL POLYPS BLENDTOME Electrosurgical Unit

Many authorities consider fulguration of rectal or sigmoidal polyps a superior method because it minimizes hemorrhage and affords better convalescence. The BLENDTOME Portable Electrosurgical Unit provides facility for fulguration, coagulation or excision of pedunculated polyps, diffuse polyps, "multiple polyps" and other tumorous conditions of the colon. Besides for the proctologist, the BLENDTOME offers advantages for the G. P. as well as the specialist. This moderate priced portable unit equips the doctor with easier technics for biopsy, cervical conization, mass removal of various growths and numerous other surgical pro-



BIRTCHER

To: The BIRTCHER Corp., Dept. A-11-9 5087 Huntington Dr., Los Angeles 32, Calif. Please send me your free brochure on the Blendtome Portable Electrosurgical Unit.

State

have a disturbance in sugar metabolism. Certainly these people should be made to reduce before insulin is given to them. It has been my experience, and I am sure it is shared by many others, that when such individuals are made to reduce they do not require insulin at all. In short, I think the medical profession should be more alert to the subject of obesity, certainly as it concerns the diabetic or potential diabetic.

I feel that to one who is not too well acquainted with the management of diabetes this comment might be misleading in the treatment of

diabetes.

J. SHIRLEY SWEENEY, M.D. Gainesville. Tex.

¶We sent Dr. Sweeney's letter to Dr. Colwell for comment. His reply follows.

TO THE EDITORS: Dr. Sweeney is correct in assuming that my summary of the indications for protamine insulin was not intended to imply that obese diabetics should necessarily be given that insulin. The incident emphasizes the inadequacy of a summary.

What I intended to say was that when a diabetic patient requires insulin at all, the one who is obese is more likely to have a mild form of the disease which is easily controlled with small doses of protamine insulin, in contrast to the one with diabetes so severe that he is thin and requires an insulin with intermediate action such as globin insulin or one of the protamine mixtures.

I also agree with Dr. Sweeney that obese diabetics should be urged to reduce, although I would not withhold insulin from them while they are reducing if it is necessary to control their sugar metabolism. The in-



effectiveness of MEPRANE DIPROPIONATE therapy — "Effective," "Highly effective,"2 "Highly efficient." In private practice, thousands of physicians are prescribing Meprane dipropionate with equal success.

Meprane dipropionate relieves menopausal symptoms "quickly and thoroughly."3 Unpleasant reactions have been reported to be "conspicuous by their absence."1

Write for literature and sample.

Dosage: Initial therapy 1 (1 mg.) tablet t.i.d. after meals; maintenance therapy, 1 to 2 tablets daily.

Packaging: 30, 100, 500 and 1000.

St.

REED & CARNRICK - Jersey City 6, M. J. - Toronto, Ont., Canada

#### Bibliography

- 1. Sturgis, S. H.: Am, J. Obst. & Gyn. 53:678, April 1947.
- 2. Coulton, D. and Sewall, C. W.: Am. J. Obst. & Gyn. 56:541-548, Sept. 1948.
- 3. Lin, H. A. C.: Am. J. Obst. & Gyn. 54:296, August 1947.





sulin can always be stopped after weight reduction if it is then unnecessary.

ARTHUR R. COLWELL, M.D.

Evanston, Ill.

## **About That Diagnostix Collection**

TO THE EDITORS: Kindly reserve my copy of the 1949 series of 24 Diagnostix. Thank you in advance.

FELIX OTTAVIANO, M.D.

Oneida, N.Y.

► TO THE EDITORS: Please reserve my copy of Diagnostix Collection. Thank you for your cooperation.

MAXWELL J. ANTELL, M.D.

Bridgeport, Conn.

¶To Drs. Ottaviano and Antell, and to all other readers who obtained the impression that the Diagnostix Collection was to be a separate volume, our apologies. The reason for the misunderstanding was the following notice which appeared on page 88 of the September 15th issue

#### DIAGNOSTIX COLLECTION

The 1949 series of 24 Diagnostix will be included in the Modern Medicine Annual—1950. Reserve your copy now.

What we meant to say in the last sentence of the notice was "Reserve your copy of the MODERN MEDICINE ANNUAL—1950 now." The Diagnostix Collection is a part of the ANNUAL and is not available separately.—Ed.

#### **Helpful Hints**

TO THE EDITORS: I enjoy your magazine and feel it has many helpful hints.

SAMUEL CISSEL, M.D.

York, Pa.



"Hello, Mr. Smith. How's the eczema today?"

## **Mixup Straightened Out**

TO THE EDITORS: I would like to call your attention to an error in the heading of the article in regard to auricular fibrillation in a normal heart that appeared in the September 15 issue of *Modern Medicine*, p. 47. The article is from the Lahey Clinic in Boston and not from the Mayo Clinic in Rochester, Minn. Dr. Hugh Hanson is now a fellow at the Mayo Clinic, but the report was written while he was a fellow at this institution.

DAVID I. RUTLEDGE, M.D.

Boston

TO THE EDITORS: Thank you for reviewing our article on "Auricular Fibrillation in Normal Hearts." The material for this article and the writing thereof came from the department of internal medicine of the Lahey Clinic.

HUGH H. HANSON, M.D.

Rochester, Minn.

¶Inadvertently we placed both authors of this article, Dr. Rutledge and Dr. Hanson, at the Mayo Clinic. Dr. Rutledge is with the Lahey Clinic. So, too, was Dr. Hanson at the time the article was written, but he has since become affiliated with the Mayo Clinic.—Ed.

# For Low Sodium, High Protein Diets

For patients who require diets low in sodium, high in protein, many physicians recommend a regime of low-sodium, high-protein main dishes, salads and desserts made with Knox Gelatine.

Knox Gelatine makes possible a simple, basic method of food preparation—for a large variety of dishes—bland, easily digested and extremely appetizing. Your patients will find suitable recipes enclosed in each package.

Knox Gelatine is not like the readyflavored gelatin dessert powders with their high sodium (and sugar) content. Knox is all gelatine—of high quality. It is all protein—no sugar, no acid, very low in sodium.

## Free Dietary Literature

A series of special booklets devoted to menus and recipes for prescribed diets are yours for the asking. Address Knox Gelatine, Dept. R-19. Johnstown, N. Y.



#### Hot Water vs. Refrigeration

TO THE EDITORS: While practicing in Arabia, I encountered three times in homes where conditions were quite primitive such severe postpartum hemorrhages that the patients' lives were endangered.

Besides elevating the legs and giving pituitrin, I gave a douche of water as hot as could be tolerated. A teaspoonful of tincture of iodine was put in the water. An ordinary enema can with a sterile nozzle was used.

In all 3 cases this heroic and unscientific measure did the trick. There was no more bleeding and the patients lived.

I adopted the measure at the suggestion of my Indian nurses who were trained as midwives and had observed its effectiveness in India.

I would not think of using this measure before trying the usual methods of control, as patients do not enjoy it. In the cases mentioned, it was a last resort—but bleeding stopped.

You may send this to Dr. Frederick M. Allen if you like. I noticed his letter about the use of refrigeration in obstetric crises in your Medical Forum recently (Sept. 15, 1949, p. 78). I'm a little doubtful whether he or most doctors will think very highly of my method, but it might conceivably save a life if some doctor caught in a home is at his wits' end to save a life, as I was.

NEOSKOLFTA TIFFANY, M.D. Inglewood, Calif.

## Hysteria in Children

TO THE EDITORS: The term hysteria may be used in either a general or precise manner. It has been used loosely to describe any type of emotional reaction—"hysterical attack." However, in its more precise form



# complete topical treatment for middle and external ear infections

- High Antibacterial Potency—high concentration of sulfa-urea at site of infection.
- Chemical Debridement—infection site rapidly cleansed—odors reduced, and waste material removed.
- 3. Analgesic and Antipruritic—pain and itching relieved by chlorobutanol.
- 4. Fungicidal Action—common fungous pathogens inhibited.
- 5. Hygroscopic—excess moisture absorbed, decongestive action.

White's Otomide is a stable solution of 5% Sulfanilamide, 10% Carbamide (Urea) and 3% Anhydrous Chlorobutanol in glycerin of high hygroscopic activity.

Supplied in dropper bottles of  $V_2$ -fluid ounce (15 cc.)



WHITE LABORATORIES, INC., Pharmaceutical Manufacturers, Newark 7, N. J.



Microthann RADAR DIATHERMY

Check for TV interference before you buy any diathermy equipment. FCC approval does not guarantee freedom from interference or from restriction if interference develops.

Microtherm employs radar frequencies way above the television wave range — no interference.

Ask your dealer to give you a demonstration or write for Bulletin DL-MED601.

## ADVANTAGES OF RAYTHEON MICROTHERM

- Penetrating energy for deep heating
- 2. A desirable temperature ratio of fat to vascular tissue
- 3. Effective production of active hyperemia
- Desirable relationship between cutaneous and muscle temperature
- 5. Predetermined and reproducible dosage no tuning
- Controlled application over large and small areas
- No contact between patient and directors, Shock-proof





Approved by the F. C. C. Certificate No. D-477
Underwriters' Laboratories

# RAYTHEON MANUFACTURING CO. Power Tube Division

Waltham 54, Massachusetts

it is usually considered to represent a conversion phenomenon. In this mechanism the individual's conflict is converted or changed into a physical symptom, usually with the loss of the normal anxiety and tension that accompany such a conflict.

Such patients may have a paralysis of the hand that is accepted with other types of psychoneurotic reactions in which the individual develops physical symptoms but retains the anxiety and tension associated with his problems. In these neurotic reactions he may have disturbances of heart, bladder, or bowel due to the normal psychosomatic reaction associated with such moods as anger, fear, or anxiety.

A true hysteria of the type usually found in adults and described above is a somewhat rare occurrence in childhood. In many instances the whole mechanism appears very close to consciousness and one often suspects that the child is deliberately putting on these symptoms in order to solve his problems.

Such symptoms are of value to the child, usually in one of two ways—by enabling him either to avoid an unpleasant situation or to obtain attention, affection, and increased security. It may well be that the basic mechanism is the same in both the child and the adult and that the difference in the final picture depends upon the emotional maturation of the two individuals, their life experiences, and their intellectual levels.

Hysterical reactions in children, while arising in various situations, are most commonly seen in relation to a school situation, when the child is having difficulty with the work and is falling behind his classmates. The

(Continued on page 28)



in the control of
ANXIETY STATES - NERVOUS TENSION
SMOOTH MUSCLE SPASM

PHENOBARBITAL gr. 1/4 (16 mgm.)

PHENOBARBITAL gr. 1/4 (16 mgm.)

BELLADONNA ALKALOIDS 0.159 mgm

BELLADONNA atropine, scopolamine in scopolamine, aropine, scopolamine in equivalent apuniform proportion and equivalent apuniform proportion and equivalent approximately to Tr. Belladonna 8 minims.)



# BELBARB

Available in

APSULES . ELIXIR



CHARLES C. HASKELL AND COMPANY, INC.

......

now, better relief with..

Robitussin

rationally formulated

clinically proved



a distinctive, new, non-narcotic

# antitussive-expectorant

At last, something really new in cough syrups . . . something completely rational . . . clinically sound . . . Robitussin 'Robins'.

Robitussin employs glyceryl guaiacolate and desoxyephedrine hydrochloride, in a palatable aromatic syrup vehicle.

Glyceryl guaiacolate has proven an effective aid to expectoration, and a cough ameliorator with prolonged action, through its increase in and thinning of respiratory tract fluid; <sup>1,2,3</sup> yet it has no ill effect upon digestion.

Desoxyephedrine's sympathomimetic action is also well recognized: <sup>4,5,6</sup> by relaxing spasm of the bronchial musculature and helping maintain normal respiratory smooth muscle tone, it greatly minimizes the provocation of cough from spasm.<sup>3</sup> At the same time it affords relief from psychic depression or a feeling of fatigue.

The syrupy vehicle, with its aromatic volatile oils, has a local demulcent effect. Furthermore, it assures patient cooperation by providing a base which makes Robitussin one of the most palatable of all antitussive-expectorants.

You will find Robitussin 'Robins' an exceptionally efficient, safe, therapeutic tool in the management of cough — for both adults and children.

DOSAGE: Children: one-half to one teaspoonful, according to age, three or more times daily. Adults: one or two teaspoonfuls, as necessary every two to three hours.

SUPPLIED: Pint and gallon battles.

REFERENCESs 1. Connell, W. F. et al: Conadian Med. Assoc.
J., 42:220, 1940, 2. Perry W. F. and Beyd, E. Mr. J. Pharm.
Exper. Ther., 73:65, 1941, 3. Stevens, M. E. et al: Canadian
Med. Assoc. J., 48:124, 1942, 4. Foltz, E. E. et al: Canadian
Med. Assoc. J., 48:124, 1942, 4. Foltz, E. et al: 51; J. Leb. Clin.
Med. 28:603, 1943, 5. Gruhom, S. E. Ind. Eng. Chem., Ind. Ed.
37:149, 1943, 6. Schulz; F. and Deckner, S.: Klin. Wecheskr.,
21:674, 1942.

A. H. ROBINS COMPANY, INC., RICHMOND 20, VIRGINIA

Ethical Pharmaceuticals of Merit since 1878

## Formula

Each 5 cc. (1 teospoonful) of Robitussia contoins Glycaryl Gualacolate, 100 mg. Dosaxyaphadrina Hydrochlaride, 1 mg. In a palatable areasetic syrup.

# Robitussin

for rational cough management







Assurance of this comes from the fact that "Viso" performance not only conforms with all A.M.A. requirements, but EXCEEDS many of them, notably those concerned with the faithfulness of reproducing wave forms. (Proof by technical data gladly furnished.) Viso-Cardiette recording characteristics even meet most research needs!

Also, recording is inkless, and in standard rectangular coordinates . . . no curvature of complexes or time lines, no negative time intervals. The baseline is steady, and free from "AC". "Viso" recording paper is 6cm. wide with a ruled area of 5 cm. width, is permanent, non-fading, and can be handled freely, filed easily.

The coupon below will bring you, without obligation, a new descriptive folder which describes more in detail the clinical advantages of the "Viso" record.

	» S	AN	B	0	R	1		C			C	A	1	Į.	S	GI
Please descrip	send	me Viso	-C	vit are	ho lie	ut	e	b	lig	ga de	ti	01	1	n	e	w
Name .		++1													×	
Street					. ,											
City &	State											11				

hysterical reaction to such a situation is usually one that will solve the problem: paralysis of the writing hand, a disturbance of vision which keeps him from seeing the board, a disturbance of hearing which prevents him from following the teacher, or abdominal complaints and pains which are usually most evident in the early morning and prevent him from attending school.

When the hysteria is secondary to a disturbance in the home situation, the pattern is more likely to be varied, but is usually one that enables the child to obtain more attention and affection from the members of the household or elevates him into a preferential position compared to his

brothers and sisters.

The treatment of hysteria requires a full explanation of the cause of the symptoms to the parents. Therapy must also be directed toward correcting the basic situation.

If the difficulty is in school, it may mean less pressure by the teacher. extra coaching, or placement in a lower grade or an opportunity class. If the origin of the symptoms lies in the home, which incidentally is a much more difficult situation to control, an attempt must be made to improve the relation between the parents and the patient and the patient and his siblings. This must be worked out on a more wholesome and satisfactory basis, one that will lead to an eventual sense of acceptance and security on the part of the child. In addition to this, one should see that the child is not allowed to escape situations through the utilization of such symptoms nor obtain excessive attention or prestige through such symptoms.

W. A. HAWKE, M.D.

Toronto



'Dexedrine' Sulfate can help restore mental alertness and optimism, dispel psychogenic fatigue and thus ''make life worth living.'' Smith, Kline & French Laboratories, Philadelphia

# Dexedrine\* sulfate tablets - elixir

\*T.M. Reg. U. S. Pat. Off. for dextro-amphetamine sulfate, S.K.F.

the anti-depressant of choice

# new convenience new flexibility in dosage new all-around usefulness



## Poly-Vi-Sol

Each 0.6 cc. supplies:

Vitamin A 5000 USP units
Vitamin D 1000 USP units
Vitamin D 1000 USP units
Ascorbic Acid 50.0 mg.
Thiamine 1.0 mg.

Riboflavin 0.8 mg. Niacinamide 5.0 mg. Tri-Vi-Sol

Each 0.6 cc. supplies:

Vitamin A 5000 USP units
Vitamin D 1000 USP units
Ascorbic Acid 50 mg.

Ce-Vi-Sol

Each 0.5 cc. supplies: Ascorbic Acid 50 mg.

# 3 new water-soluble liquid vitamin preparations

Each of these preparations is ideally suited for routine prophylactic or therapeutic vitamin supplementation for infants and children as well as adults.

Water-soluble, pleasant tasting, they can be stirred into the infant's formula, or into fruit juice, milk, or other fluid; mixed into cereals, puddings, etc.; or incorporated in mixtures for tube feeding.

Each is scientifically formulated and ethically marketed. They are supplied in 15 and 50 cc. bottles, with an appropriately calibrated dropper.

MEAD'S

MEAD JOHNSON & CO. EVANSVILLE 21, IND., U.S. A.

# HORMOTONE "T"

## **Enterosol Coated Tablets**

Estrogenic Hormones and Thyroid Orally Administered

Acts directly upon the endometrium inducing hyperplasia of the uterine mucosa.

For the treatment of ovarian hypofunction: amenorrhea, certain types of dysmenorrhea, hypomenorrhea, oligomenorrhea and menopausal disorders.

Each tablet contains, 1,000 international units of biologically assayed and standardized estrogenic hormones combined with 1/10 grain thyroid.

Also in tablets containing 5,000 international units each.

Bottles containing 20, 100 and 500 Enterosol Coated tablets.



G. W. Carnrick Co. 20 Mt. Pleasant Ave.

Newark New Jersey

## Therapy for Aphthous Stomatitis

TO THE EDITORS: I fully agree with Dr. J. M. McMahon's conclusion relative to the psychosomatic etiology of certain cases of aphthous stomatitis, that terribly disturbing condition about which so little mention has been made in the nation's medical literature (Sept. 15, 1949, p. 18). Another discouraging phase of this condition has been the long list of therapeutic suggestions, few of which are even worth trying.

In four decades of practice, I have encountered few conditions so responsible for localized poignant pain in the region where these oval-shaped, pearly tinted, ulcer-like sloughs occur. I have had patients state emphatically that if they had to choose between the intense pain of a furuncle or carbuncle and the patch of an aphthous stomatitis, they would much prefer the former. I doubt very much whether there are many conditions in the practice of medicine which find a physician so thoroughly helpless.

Recently I encountered a case of a very persistent type which was frequently recurrent for over a year. My reaction, I imagine, was like most physicians' when called upon to treat this disturbance. One of the ulcers seemed to be unduly irritated, and since the patient was a middle-aged individual, I called into consultation a competent specialist in cancer and allied conditions, who nevertheless does not overlook his general therapeutics.

His decision was to place the patient on large doses of a high potency yeast powder. Progress has been somewhat slow but definite, and it does appear at this time that the condition will clear up.

WILLIAM HARVEY THALER, M.D. Long Beach, Calif.

# how to get rid of undesirable tenants

In Vaginitis
and Carvicitis
hardy indeed is the
trichamonal or other
infective organism
which can survive
the vaginal
environment
created by

# westhiazo

Safe, dainty, easy-to-use westhioxole vaginal rapidly produces . . .

- a vaginal acidity untenable to most pathogenic organisms.
- Speedy control of discharge, itching, foul odor, and other distress.
- More rapid recovery by elimination of secondary as well as primary infection; recovery in vaginitis averages 2 to 7 weeks; in cervicitis 3 weeks.

westhiozole vaginal jelly
contains 10% SULFATHIAZOLE,
4% UREA, 3% LACTIC ACID,
1% ACETIC ACID in a
polyethylene glycol base

samples? literature? please write to
WESTWOOD PHARMACEUTICALS, Dept. MM
468 Dewitt St., Buffalo 13, N. Y.
division of Foster-Milburn Co.



# NOW! ONE NEW material replaces 4

. SHEET WADDING . CREPE PAPER . WOOL FELT

. STOCKINETTE IN ORTHOPEDIC CASTS

The First and Only True, Non-Woven, All-Cotton Felt

# Curity ORTHOPEDIC BANDAGE

There is no other material like it! This revolutionary bandage material, a new non-woven, all-cotton felt long believed impossible to manufacture, is considered indispen-

sable by surgeons who have used it experimentally. The fabric is an entirely new material—developed after years of testing and research by The Kendall Company.

#### MANY ADVANTAGES FOR USERS

- 1. Strength-holds together, wet or dry.
- 2. Conformability—adjusts to body contours, no wrinkling.
- 3. "Cling" or Cohesiveness-adheres to itself, cannot delaminate.
- 4. Elasticity provides valuable support; not just bulk.
- 5. Ease of Application—simple, quick, lint-free.
- 6. Smooth, yet Non-skid-ideal surface over which to apply plaster.
- 7. Durability—does not wad or bunch up under cast.
- 8. Absorbency—skin condition is protected...does away with many skin problems.
- 9. Porous-feels snug, yet "live"-air can circulate.
- 10. Easy to Cut—cast removal is an easy, clean job.
- 11. Non-shrinkage does not get tighter after moistening.
- 12. No Waste odd lengths can be used for padding, stay where placed.

#### OTHER SUGGESTED USES:

- rectal dressings eye pads fluffs
- colostomy care
   bandaging
- sponging
- vaseline dressings

A product of

#### (BAUER & BLACK)

Division of The Kendall Company, Chicago 16

Ask Your Curity Representative to Demonstrate



# Washington Letter

## Physician's Interest in National School Health Program

Before long, most physicians will come into direct contact with provisions of the national school health law. This program authorizes federal assistance to states by a formula that allows a larger per-pupil allowance for the low-income sections. A total of \$35,000,000 is provided for these grants.

The basic idea is to insure 'hat no American child shall come to adult life with physical or mental defects or conditions which can be prevented or corrected at an early age. To this end, the program provides aid to the states for "prevention, diagnosis and

end, the program provides aid to the states for "prevention, diagnosis and treatment of physical and mental defects and conditions of all school chil-

dren, with special reference to the correction of defects and conditions likely to interfere with the normal growth and development and educational progress of children."

The plan originally was attached to the general federal aid to education bill. Sen. Elbert Thomas of Utah, chairman of the Senate committee, made it a separate bill early in the last session of Congress. He anticipated

the possibility of a controversy over aid to private and religious schools and decided that, on health, this argument should not be allowed to arise. As a result, all children will receive the same attention, whether public, private, or parochial pupils.


For the private physician, several points in the law deserve careful attention when funds become available on the local level.


1 The law requires that medical and dental examinations be provided at the schools, as long as any federal funds are available for the purpose.

(Continued on page 41)



B-D Mesents





new precision new sensitivity new visibility new serviceability

Look for the red dot ● when you
buy an ANEROID MANOMETER
. . . it identifies the Jeweled · Bearing
that gives greater precision,
performance and permanence.

### check these advantages:

Jeweled · Bearing reduces friction, maintains accuracy, lengthens life.

Uniformly-spaced scale graduations provide greater visibility; registers from 0 to 314 mm.

Long-travel beryllium copper bellows assure dependability and years of service.

Rugged construction . . . Luer slip connection permits instrument to be removed easily from inflation system for easier deflation, for interruption during exercise test; for use on several cuffs in clinic practice, and for use on different size cuffs. The manometer can be left

YALE ANEROID MANOMETER

in blackt or sorgical gray.

Indeptivitely government against all defects in material or specimenship.


Supplied in black platfin entra-roomy saveleys you have been a platfin as the facts of the same of the sam

### and the B-D

## SECURITY CUFF

A hook-and-eye cuff that is simplicity itself. Takes but a few seconds to apply; no waste motion, no loose ends; no bulging or herniation; gives positive, secure attachment to instrument; offers the flexibility of 20 adjustments.





The B-D Security Cuff is regularly supplied with B-D Manometers.

See the new B-D YALE ANEROID MANOMETER

Jeweled · Bearing with the B-D Security Coff at
your surgical dealer's.



BECTON, DICKINSON AND COMPANY, RUTHERFORD, N. J.

2 The law provides for the utilization, insofar as possible, of "the qualified health, medical, dental and hospital facilities already established in each community, and if no state law forbids, with special reference to utilizing the services of the family physician."

For practical purposes, this means that no physicians or dentists may be brought into a community to handle the program, if adequate professional manpower is available in the community.



Furthermore, the law states that representatives of professional associations shall be members of a state committee to advise the state health department in administration of the act. Active and well-informed representatives on these boards should be

able to protect the interests of the profession. However, if these delegates are not sufficiently active, or are insufficiently informed, difficulties may arise.

3 The law requires that federal funds, as long as they are available, be used to pay for treatment of defects shown in the examination "whenever the parents of such children are unable to provide such treatment." Application of this last phrase deserves careful attention by the family physician. Each state will be allowed to set its own criteria for determination of when a family is "unable to provide" treatment. In the case of a low-income family, this point might be of importance to the physician.

A long-debated point is whether states should be allowed to let the plan cover treatments for all children, without a test of the family's ability to pay. As this is written, in

(Continued on page 146)

# Questions & Answers

All questions received will be answered by letter directed to the petitioner; questions chosen for publication will appear with the physician's name deleted. Address all inquiries to the Editorial Department, MODERN MEDICINE, 84 South Tenth Street, Minneapolis 3, Minnesota.

QUESTION: A man thirty-five years old had a compound fracture of his left tibia and fibula about fifteen years ago. The leg healed well, but the left calf is now 21/4 in. larger than the right. Kindly advise the proper procedure.

M.D., New Jersey

ANSWER: By Consultant in Orthopedics. The difference in size of the two calves probably cannot be adjusted without surgery; it is doubtful that overexercise of the smaller calf would provide permanent increase in bulk of muscle. If no symptoms other than discrepancy in circumference exist, it would be best to accept the circumstance.

QUESTION: I have a young patient in her second month of pregnancy who has severe leg cramps at night. Can you tell me what causes the cramps and what to do to prevent them?

M.D., New York

ANSWER: By Consultant in Obstetrics. The cause of leg cramps during pregnancy is not entirely clear. The most popular theory is that calcium deficiency leads to tetanic contractions of the muscles of the lower extremity. These occur during sleep because the circulation becomes stagnant in the absence of muscular contraction. Trousseau's sign is based on similar conditions in the upper extremity except that the vascular stasis is artifi-

cially produced. The advocates of the above theory attempt to increase the calcium intake to combat the cramps. Others have suggested thiamine deficiency as contributory and give large doses of vitamin B<sub>1</sub>. Fortunately, massage and walking promptly relieve most patients.

QUESTION: Could you give me references on the use of urine injections in the treatment of urticaria?

M.D., New Mexico

ANSWER: By Consultant in Dermatology. The use of protein extracts from urine in treatment of urticaria was suggested in England some years ago but, as far as I know, the treatment has not been followed there and certainly not in this country. The following references are suggested:

 Herz, K: Ueber eigenharnbehandlung. München. med. Wchnschr. 78:398-400, 1081.

2] Gutiérrez, P., and Dionisio, S. A. Autourotherapy in urticaria. Acta med., Philippina 2:427-433, 1941.

3] Jausion, H., and Paléologue. Une nouvelle méthode de désensibilisation; l'auto-ouro-thérapie dans la cure de l'eczéma. Bull. soc. franç. de dermat. et syph. 36:115-118, 1929.

et syph. 36:115-118, 1929.

4] Jausion, H., Carrot, E., and Gibert, A. Un cas d'eczéma solaire spontané. Sa brusque généralisation après application de pommade. Sa guérison l'auto-ouro-thérapie. Bull. Soc. franç. de dermat. et syph. 38:9-14, 1931.

IN DAYTIME SEDATION



MENOPAUSAL HYSTERIA



HYPERTENSION



**OBSTETRICAL HYPNOSIS** 



PREOPERATIVE APPREHENSION



INSOMNIA



# **butisol** sodium

BRAND OF BUTABARBITAL SODIUM

intermediate sedative

Butisol Sodium produces "a relatively mild and more continuous depression than can be obtained with the shorter-acting barbiturates, yet its action is less prolonged than with barbital or phenobarbital."

With proper regulation of dosage, there is no cumulative action and a minimum of lethargy and "hang-over."

Sedation is sustained for approximately five to six hours—without sharp peaks of effect—thus providing a most useful sedativehypnotic in a wide range of clinical indications.

DOSAGE FORMS: Elixir Butisol Sodium 0.2 Gm. (3 gr.) per fl. os. Also Capsules and Tablets, 0.1 Gm. (1½ gr.); Tablets, 15 mg. (½ gr.), 30 mg. (½ gr.) and 30 mg. (½ gr.). Caution: Use only as directed.

Elizir butisol sodium

Its bright, green color and refreshing flavor appeal to all; excellent prescription vehicle. Clinical samples on request.



1. J.A.M.A. 135:224 (Sept. 27) 1947.

### McNEIL

LABORATORIES, INC., PHILADELPHIA 32, PA.



# PERTUSSIN increases the RTF\* which is the ABC of Cough relief

—in acute and chronic bronchitis and paroxysms of bronchial asthma ...in whooping cough, dry catarrhal coughs and smoker's cough. PERTUSSIN increases the Respiratory Tract Fluid which is the key to its effectiveness in relieving such coughs.

PERTUSSIN therapy is simple but fundamental. It lends a helping hand by the practical device of assisting nature to work in its own defense. No wonder PERTUSSIN has been in successful use for over thirty years!

Entirely free from opiates, creosote and chloroform, PERTUSSIN is well tolerated—without undesirable side action—by children and adults alike, and is pleasant to take.

\*Respiratory Tract Fluid

## PERTUSSIN

For Children, Adults and the Aged

SEECK & KADE, INC. NEW YOLK 13, N. Y. 5] Johnston, H. A. Autodesensitization of allergic conditions. California & West. Med. 4:307-309, 1934.

6] Nicastro, A., and Petronici, G. L'autouroterapia in dermatologia. Riv. san.

siciliana 26:867-872, 1938.
7] Tereshkovitch, V. I., and Tokareff, A.P. Autourinotherapy in skin diseases (in Russian). Vrach. gaz. 34: 1583-1586, 1930.

### QUESTION: Can you recommend treatment other than x-ray for keloids? M.D., Georgia

ANSWER: By Consultant in Dermatology. The best treatment for keloids is probably radiotherapy by x-rays or radium. Nevertheless some cases are resistant and other measures may be applied. Occasionally a satisfactory result is obtained with application of solid carbon dioxide. Small lesions can sometimes be treated successfully by shaving with a sharp blade and then light electrofulguration.

Keloidal reaction may occur in one scar but not in others in the same individual. Under this circumstance excision may be tried and the site watched closely for evidence of new keloidal development. If the area has not previously had large amounts of radiotherapy, that treatment may then be given in order to prevent further keloidal change.

## QUESTION: When is the sex of the fetus determined? M.D., New York

ANSWER: By Consultant in Genetics. Sex determination probably occurs at the time of conception and depends upon the type of chromosomes carried by the sperm. Secondary sexual characteristics may be influenced by maternal hormone production. Sexual development has been influenced by extrinsic hormone administration in animals but not in human beings.

### "no signs of renal irritation

### were encountered"

A mixture of 3 sulfonamides offers a "markedly reduced" incidence of crystalluria and renal irritation . . . and "is significantly less toxic" than a single sulfonamide, or a mixture of two.

"Therapeutic results"—in one thousand unselected patients with acute systemic infections treated with sulfonamide combinations—"were uniformly satisfactory and conspicuous in many instances by the speed of clinical improvement. Crystalluria was infrequent . . . No signs of renal irritation were encountered. The incidence of allergic reactions also appeared decreased."

1. Lehr, D.: Presented at The Scientific Exhibit, American Medical Association, June 21-25, 1948.

and tableta

safer,

more effective sulfonamide therapy

Each 5 cc. of syrup (approx. one teaspoonful) ... or each tablet contains 71/2 grains of sulfa compound:

Sulfadiazine	0.162 Gm. (2½ gr.)
Sulfamerazine	0.162 Gm. (2½ gr.)
Sulfathiazole	0.162 Gm. (21/2 gr.)

Tri-Sulfanyl Syrup also contains sodium citrate 0.375 Gm. (5.8 gr.) in a pectinized, vanilla flavored base.

Samples on request.

casimir funk laboratories, inc. affiliate of u. s. vitamin corporation 250 E. 43rd St., New York 17, N.Y.

Ideal For Premature, Normal Babies



# Finish EVENFLO Bottles Better!

Babies sometimes go to sleep before emptying their bottles because their limited strength is exhausted trying to get food thru a hard, stiff

or collapsed nipple.

Not so with Evenflo. The Evenflo Nipple is soft and pliable throughout so that baby nurses by compression as well as suction, similar to breast feeding. The twin air valves keep air pressure in bottle constant, allowing formula to be withdrawn easily when nursed. Because they nurse Evenflo in comfort, babies get more benefit from their food and make better gains in weight.

It's this smooth nursing action of Evenflo's patented twin-valve nipple that has made it—

America's Most Popular Nurser



QUESTION: For about eight years a woman twenty-eight years old has had a temperature fluctuating daily from 98 to 101° and often to 103°. Her only other complaints are morning fatigue, headaches in the morning and evening, and an ache in her left arm. All laboratory data are negative. Recently she took aureomycin, 16 capsules, 250 mg. daily, for two days. Temperature was normal for a day or two but now rises daily to 101°. She has, however, felt better since taking aureomycin. Can you suggest a diagnosis?

M.D., California

ANSWER: By Consultant in Internal Medicine. The temperature variations described, if known by accurate observations for a period as long as eight years, are probably normal for the individual. Similar cases have been reported, notably by Hobart Reimann. Morning fatigue and evening headache are usually psychogenic symptoms and are not related to changes of body temperature or to a particular disease.

QUESTION: Could you give me the technic, or reference, for artificial insemination along with precautions and contraindications? I have a married patient, age twenty-nine, who is very anxious to become pregnant.

M.D., Kansas

ANSWER: By Consultant in Obstetrics. Excellent results have recently been reported by Douglas P. Murphy, M.D., and Edmond J. Farris, Ph.D., utilizing the rat ovulation test to detect the day of ovulation. Their technic is reported in J.A.M.A. 138:13 (Sept. 4) 1948. The basal temperature record may also be helpful for timing purposes. Evidence of infection in either the semen or the cervix is a contraindication to artificial insemination. Obviously, complete sterility studies must precede insemination and reveal no serious obstacle to conception.

description

Tyrozers are pleasantly flavored, pink lozenges, each containing 1 mg. of antibiotic tyrothricin, and 5 mg. of soothing, analgesic benzocaine.

## indications

Topical treatment of sore throat associated with colds, hay fever, and other allergies, or resulting from chemical irritants or vocal strain also postsurgical care of the pharynx.



Tyrozets quickly relieve the pain of sore throats (benzocaine), and help suppress local infections due to a wide range of gram-positive organisms (tyrothricin).

# Tyrozets

Antibiotic-Anesthetic Throat Lozenges

Supplied in unbreakable, amber-plastic vials of 12 lozenges. Sharp & Dohme, Philadelphia 1, Pa.



### What Would You Say?

Twice a month we will select a caption for this cartoon from those sent in by our readers and send the author \$5. This caption was written by James L. Tullis, M.D.

Mail your caption to The Cartoon Editor. MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.

Brookline, Mass.



With this disease you have to give up only smoking and drinking."



re

fc

fo

de



# OF PAIN AND LESIONS OF HERPES

DESCRIPTION: Protamide is a sterile, aqueous colloidal solution of a specially processed proteolytic enzyme, for the maximum relief of nerve root pains of Herpes Zoster and Tabes Dorsalis.

CLINICAL RESULTS: Highly gratifying clinical results have been obtained with the use of Protamide (Sherman) in the treatment of the extremely resistant herpes syndrome. Pain has been relieved in the great majority of herpes cases within four to forty-eight hours and lesions have healed in ten days or less—regardless of the particular nerve roots involved. Complete clinical data may be obtained by writing for the Protamide literature on Herpes Zoster and a recent reprint on Protamide for Tabes Dorsalis.

DOSAGE: In Herpes Zoster the recommended dosage is 1.3 cc of Protamide intramuscularly each day from two to four days. No contraindications or incompatability have been reported to date. All Protamide is clinically tested for positive results. Can be stored at room temperature without loss of potency.

# HERPES ZOSTER

ALSO CLINICALLY PROVED
FOR THE LIGHTNING PAINS
AND ATAXIA
OF TABES DORSALIS

\* REGISTE

SHERMAN LABORATORIES

G. H. Sherman, M. D., Founder

BIOLOGICALS • PHARMACEUTICALS

DETROIT 15, MICHIGAN

FD U. S. TRADE MARK

## Forensic Medicine

COMPILED BY ARTHUR L. H. STREET, L.L.B.

PROBLEM: [1] When licensed as an obstetrician and chiropractor, is one under the same legal responsibility as a general medical practitioner in the treatment of postparturitional conditions? [2] Is the liability for negligence in diagnosis and treatment diminished by the temporary entrance into the case of a regular physician for the purpose of delivering a stillborn child? [3] In such cases, does the fact that defendant is a chiropractor and midwife disqualify a regular medical practitioner from testifying as an expert as to the sufficiency of the diagnosis and treatment, the witness knowing what a midwife actually does or should do under given conditions?

COURT'S ANSWERS: [1] Yes. [2] Yes. [3] No.

This decision was made by the Illinois Appellate Court, First District. Third Division (84 N. E. 2d 843).

PROBLEM: A sprained ankle sustained by a railway passenger on a station platform rendered her a permanent cripple. Was she necessarily debarred from collecting damages from the railway company to the extent that her injury was aggravated by her failure to consult a physician?

#### COURT'S ANSWER: No.

The Kansas City, Mo., Court of Appeals noted that the injured woman, apparently of moderate means, first thought that the sprain would yield to home treatment, although it was painful from the beginning. It

was for the jury to say whether she was grossly neglectful in failing to consult a physician and in relying on "her own homely skill.... In many poor families physicians are not called except in extreme cases. Ordinary injuries, such as cuts, bruises, burns, and sprains are given none but home treatment. Courts should be slow to condemn as contrary to rules of law the ordinary practices and usages of the large class of people who must practice the closest economy to make ends meet" (140 S. W. 602).

PROBLEM: In a suit against a surgeon for an allegedly unauthorized subtotal hysterectomy, performed in Ventura, Calif., did the trial judge err in refusing to permit a Los Angeles surgeon, called to testify for plaintiff, to state that there was no emergency justifying the operation?

### COURT'S ANSWER: No.

The California District Court of Appeal, Second District, Division 2, decided that the fact that Ventura lies in a rural community sixty miles away from Los Angeles, which possesses "many large modern hospitals," disqualified a Los Angeles surgeon from testifying to the correctness of practice in Ventura. Likewise, it was proper to exclude similar testimony of a doctor who had practiced in New York City and Los Angeles. It was a technical error to refuse to permit a third



in arthritis

• unlike gold and other drugs

# safe detoxifying colloidal sulfur

- relieves pain
- reduces joint swelling
- produces clinical remission even in the most severe cases

Sulphocol colloidal sulfur compound



Sulphocol Capsules (5 gr.) 1 or 2 after meals. Bottles of 100. Sulphocol Sol (parenteral), 25 cc. vials; 12 and 100-2 cc. vials. 1/4 to 1/2 cc. intramuscularly at 3 to 7 day intervals, gradually increased to 3 cc. Write for literature and samples of Sulphocol Capsules.

A Product of the Mulford Colloid Labora THE NATIONAL DRUG COMPANY, PHILADELPHIA 44, PA.



Manufacturers of Pharmaceutical, Piological and Biochemical Products for the Medical Profession



Now Hospitals Are "Banks," Too!

Doc Simpson was saying, "Hospitals are building up 'bone banks' that work just like blood banks. When bone is needed, the surgeon takes one from a refrigerator, cuts it to the right shape and simply splices it in."

"You doctors are sure making progress," I says, "but tell me, are any of the patients fussy about whose bone they're getting?"

"No sir!" replies Doc. "No more than they worry about whose blood they get. No one yet asked for a bone from a man who went to the same school or church he did."

From where I sit, it would be a better world if we were half as willing to accept other people's ideas and tastes, as we seem to be willing to accept their bone and blood. There'll always be differences. Some like buttermilk, others would rather have a sparkling glass of temperate beer. But underneath we're pretty much the same—deserving each other's respect and tolerance!

Joe Marsh

Copyright, 1949, United States Brewers Foundation

doctor who had practiced in Ventura County and an adjacent county to state whether the practice was the same in both counties, but the error was not prejudicial because a fourth doctor who resided in the county where the operation was performed testified for plaintiff that no emergency justifying the operation existed (208 Pac. 2d 68).

PROBLEM: An employee's claim for a workmen's compensation award for alleged injury to his spine was opposed by the employer because the claim was not filed within the time required by the compensation act. Was the employer debarred from asserting that defense, in the light of the following facts? The employee was treated by his own physician and did not consult the employer's doctor until he was ready to go back to work. The employer's doctor, examining the patient only to determine his fitness to return to work, made some indistinct roentgenograms and informed the workman that because of a muscular condition he was no longer fit for manual labor. Afterward a third doctor discovered that the employee had congenital back weakness and injury to the spine.

COURT'S ANWER: No, the employer was not barred from pleading delay in filing the claim.

The Utah Supreme Court decided that the employer's doctor's misdiagnosis was not fraudulent but, at most, due to lack of skill or care in reading the indistinct films. The court further decided that the patient reported to the employer's doctor only to determine his fitness to return to work and did not rely upon what the employer's doctor told him as to the nature of his injuries. The doctor who had treated him did not discover the injury, so why should the employer's doctor be regarded as having neglected to use good faith? (206 Pac. 2d 715.)

IN FUNCTIONAL CONSTIPATION
OF ALL TYPES, MUCILOSE INDUCES
PHYSIOLOGIC LAXATION BY
HELPING TO RESTORE NORMAL
RECTAL REFLEX ACTION. IT IS
A BLAND, BULK LAXATIVE
OF THE MUCILAGINOUS TYPE
THAT DOES NOT INTERFERE
WITH NORMAL DIGESTION.

A CONTRACTOR OF THE PARTY OF TH



## **UCILOSE**

for physiologic laxation

MUCILOSE FLAKES CONCENTRATED

MUCILOSE FLAKES WITH DEXTROSE (special formula)

MUCILOSE GRANULES WITH DEXTROSE (special formula)

Dose: 1 or 2 teaspoonfuls with 2 glasses of water twice daily.

Obtainable in 4 ounce and 16 ounce containers.

Winthrop Stearns ...

New York 13, N.Y.

Windsor, Ontario

# Immediate BACTERIOSTATIC LEVELS plus PROLONGED BACTERIOSTASIS

# PFIZER crystalline PROCAINE PENICILLIN 6 with buffered crystalline SODIUM PENICILLIN 6 for Aqueous Injection

A single 1 cc. 400,000 unit injection of the new Pfizer Crystalline Procaine Penicillin G with Buffered Crystalline Sodium Penicillin G for Aqueous Injection provides all the advantages of a repository penicillin plus the immediate high blood levels attained only with a fast-acting soluble penicillin. Note these outstanding advantages:

- Immediate bacteriostatic action 100,000 units of buffered sodium penicillin G provides high blood levels rapidly.
- Prolonged bacteriostasis 300,000 units of procaine penicillin G provides sustained levels in essentially all patients for 24 hours with a single 1 cc. injection.
- No pain No oil embolisms contains no oil and no wax suspended with aqueous diluents.
- 4. Easy to prepare and administer just add diluent to inverted vial and shake well. May be administered with an ordinary 20 gauge needle and with a regular syringe. Aqueous diluent makes it easy to clean equipment.

Pfizer Crystalline Procaine Penicillin G with Buffered Crystalline Sodium Penicillin G for Aqueous Injection is available to the medical profession through a number of leading pharmaceutical companies. Chas. Pfizer & Co., Inc., 630 Flushing Avenue, Brooklyn 6, N. Y.

400,000 Unit Single-dose Vial 2,000,000 Unit Multiple-dose Vial 4,000,000 Unit Multiple-dose Vial

Each cc. supplies 100,000 units of buffered sodium penicillin G and 300,000 units of procaine penicillin G when suspended with aqueous diluents.



World's Largest Producer of Antibiotics

Manufacturing Chemists Since 1849



## "It's the same thing every month!"


But It Needn't Be!

There's never a time when Anacin can bring more gratifying relief than during the menstrual period. Why not advise your patients who suffer periodic cramps to take Anacin tablets. Its dependable A-P-C formula will give fast, effective relief with a duration of action that exceeds the effectiveness of plain aspirin.

Anacin is easily available to you and your patients at all hospital pharmacies and drug stores.







## WYAMINE

Mephentermine\*Wyeth

Safe

to use

even

at night

\*N-methylphenyl-tertiary butylamine **WYAMINE**, either as the volatile base or the watersoluble sulfate, is an effective new vasoconstrictor, possessing several important advantages. It produces a local vasoconstriction approximately equal to that of ephedrine, but its action is very rarely attended by returgescence, or by nervousness, excitability or insomnia.

Four Convenient Dosage Forms:

WYAMINE INHALER

WYAMINE-PENICILLIN

CAPSULES for Preparation of Nasa

Preparation of Nasal Solution SOLUTION WYAMINE SULFATE

SOLUTION WYAMINE-TYROTHRICIN

Use with dropper or JETOMIZER®



# Special Article

## Papillary Adenocarcinoma of the Kidney

W. CALHOUN STIRLING, M.D.,\* AND J. E. ASH, M.D.+

Prepared for Modern Medicine

AMAJOR problem in reducing the high mortality rate with kidney tumors lies in the difficulty of early detection of the lesion. Moreover, even though new urologic technics are rapidly decreasing diagnostic delay, the feasibility of surgery must be carefully weighed before treatment is undertaken.

The physician should constantly bear in mind that 50% of patients die within two years after removal of renal tumors.

#### OCCURRENCE

Papillary adenocarcinoma is the most common tumor of the kidney and occurs chiefly in patients between fifty and seventy years of age, more frequently in men than in women. Patients usually die within two years from general debilitation or metastasis.

The tumor-is ordinarily solitary and located at a pole of the kidney, usually the lower. It attains great size and is frequently extremely vascular. Different portions vary greatly in appearance, being soft and yellow where the clear cells predominate and dark where there has been hemorrhage. Strands of fibrous stroma divide the cancer into lobules. There may be cysts or calcification secondary to cell necrosis.

The tumor tends to extend into the pelvis of the kidney and invade the renal vein. It may grow slowly and is often overlooked until metastasis is noted, usually in the lungs or bones.

<sup>\*</sup> Consultant to Bureau of Medicine and Surgery, Navy Department; Consultant in Urology, Walter Reed General Hospital, Washington, D. C. † Colonel, U.S.A. Retired.

As much variation may appear in the histology of the metastatic lesions as in the primary growth.

### DIAGNOSIS

The symptoms of blood in the urine, pain, and mass in the flank are seen and, when concomitant, make the diagnosis of kidney carcinoma obvious. These warning signs occur, respectively, in 50, 35, and 10% of all cases. Loss of weight and

strength occurs in 15%.

Physicians must continue to impress the laity with the grave import of the passage of blood from the urinary tract. Such an occurrence demands an immediate examination. The amount of blood may be slight, yet may constitute the only indication of the tumor for a long time. Because bleeding may be painless, intermittent, and of short duration, the patient is often lulled into a false sense of security. Sometimes blood clots block the ureter, causing colic-like pain on the affected side.

A large mass in the loin is usually a late manifestation of tumor of the kidney and indicates a poor prognosis. Blood-borne metastases may cause gastrointestinal pain, constipation, jaundice, headache, cough, and shortness of breath. Fever, leukocytosis, hypertension, and varicocele on the affected side are frequent. Renal tumors become very large, are tender on

pressure, and may not move with respiration.

When cancer of the kidney is suspected, a preliminary intravenous urogram should be done to ascertain the size, function, and appearance of the calyces and pelvis of the kidney. Then a cystoscopic study is made, including bilateral retro-

grade pyelograms and a differential function test.

Retraction, blunting, amputation, or absence of a major calyx is the most characteristic pyelographic finding and is observed in most cases. The tumor may extend toward the periphery and cause the calyx to have a "spiderleg" appearance. The number of calyces involved increases with the size of the growth, but a major calyx is usually primarily affected.

The examiner should not rely too much on the findings in a single pyelogram. We have seen 2 cases in which the outlines of the calyces and pelvis were normal in the retrograde pyelo-

gram, yet deformity was found on intravenous study.

Other evidence of deformity includes compression of the kidney pelvis so that flattening, narrowing, or obliteration is produced. Filling defects from blood clots or a tumor may be noted. The kidney is sometimes displaced laterally, superiorly, or medially. Abnormal insertion of the ureter is a very significant finding. Deformity of the ureteropelvic junction or alteration in the silhouette of the kidney is frequently observed.

By cystoscopic examination, hydronephrosis of the closed type may be difficult to differentiate from tumor, especially when the upper ureter is obstructed. Since nonrenal retroperitoneal tumor may displace the kidney considerably, renal mobility should always be calculated. There is less retraction of the calyces in polycystic kidney than in tumor. When retraction does occur, the calyces are broad, in contrast to the narrow calyces found in neoplasm.

The pyelogram in solitary cyst is often unaltered and the urine normal. The position of the kidney axis may be consider-

ably altered by the weight of the cyst.

Other lesions to be considered in differential diagnosis are pyonephrosis, solitary or polycystic kidney, and infections producing hematuria and pyelitis cystica or glandularis. Ptosis or rotation of the kidney or a double kidney may be mistaken for tumor. In 5 cases we have found prostatic hypertrophy associated with renal cancer. Tumor of the kidney should be considered even with prostatism if blood is passed.

An exploratory operation, implying nephrotomy, is not indicated, particularly if preliminary roentgen-ray therapy has been given. Some kidney tumors are so radiosensitive that irradiation will cause marked shrinkage and erroneous conclu-

sions may be made from the distorted pyelogram.

Occasionally, evidence of deformity is identified on intravenous pyelography, then proved erroneous by a retrograde pyelogram. In such instances the retrograde pyelogram should be repeated at regular intervals until the possibility of carcinoma is ruled out.

Aspiration biopsy may be performed in doubtful cases, particularly if cystic disease is a possibility. Arteriography should be considered if the surgeon is still unable to differentiate cysts and tumors of the kidney from other retroperitoneal masses.

### TREATMENT

With a large fixed tumor of short duration, the advisability of operation is questionable, especially if the patient's condition is below par. In such cases, the mass is obviously highly

malignant.

For such patients, a course of roenten-ray therapy has been used in the past. In Wilms's tumor, usually seen in the young, early surgical extirpation followed by roentgen-ray therapy is being used successfully. The size of a radiosensitive tumor may be greatly reduced by application of roentgen rays, though this modality does not make an inoperable tumor operable.

Metastasis should be sought before operation and roentgenograms made of the lungs, lumbar spine, pelvis, long bones, and skull. Metastases are more common with highly malignant tumors. The renal carcinoma has been extirpated in some cases in hope that its removal will decrease the metastatic process.

Calcification within the tumor indicates necrosis and has

little bearing on the degree of malignancy.

The approach requires special consideration in surgery for a large kidney mass. We find the transperitoneal route very satisfactory. It offers a valuable method of obtaining early and adequate exposure of the pedicle and exploration of the renal vein so that the mass can be removed with less trauma and the tumor tissue is not squeezed into the renal vein.

The lumbar route is preferable for small tumors or if the pelvis of the kidney is involved or diagnosis uncertain. If the lumbar approach is used for removal of a large tumor, resection of one or more ribs will facilitate excision.

Postoperative radiation is rarely useful if removal has been complete.



### Potassium Deficiency

J. W. Hollingsworth, M.D.\* Duke University, Durham, N.C.

YPOKALIEMIA, low blood potassium, may be a feature of several disorders of metabolism or of diseases of the kidney or bowel.

The lack of potassium interferes with normal impulse transmission across the neuromuscular junction. The consequence is muscular weakness or paralysis. The electrocardiogram may reveal flattening of T waves, depression of the ST segments, and prolongation of auriculoventricular conduction.

J. W. Hollingsworth, M.D., summarizes the important features of several conditions which are often associated with potassium deficiency.

Periodic paralysis is a complex, usually familial, metabolic disturbance, beginning as weakness in the legs and spreading to involve other, sometimes all voluntary muscles. An attack lasts a few hours to several days and is often nocturnal in onset. Fatigue, exposure to cold, and overeating may precipitate the paralysis, which is always bilateral.

The disease is probably intimately associated with disturbed carbohydrate metabolism. Administration of potassium salts completely abolishes symptoms.

Infant diarrhea, when severe, can cause a loss of potassium. Sodium ions then move intracellularly to replace \* The clinical significance of potassium deficiency. Bull. School Med. Univ. Maryland 34:1-10,

the potassium. Acidosis is thus increased. Potassium should be included in the fluid and electrolyte replacement therapy.

A recommended parenteral solution contains 2 gm. potassium chloride, 3 gm. sodium chloride, and 40 cc. molar lactate in 710 cc. of distilled water. From 80 to 150 cc. of the solution is given per kilogram of body weight daily. Potassium should not be given if urine excretion is suppressed because potassium intoxication may result.

Diabetic acidosis may be complicated by hypokaliemia. Several factors contribute to the potassium deficiency: dehydration, washing out of electrolytes by parenteral fluid therapy, and the utilization of potassium by glycogen deposition in the liver and muscles when insulin is given.

Therapy of diabetic acidosis should include total electrolyte replacement, not only sodium chloride and water.

Addison's disease treated with desoxycorticosterone is sometimes accompanied by hypokaliemia, especially when the patient is eating a low-potassium diet. The drug causes sodium retention at the expense of excessive potassium excretion.

Steatorrhea can increase loss of potassium in the feces. Unabsorbed intestinal fat combines with various bases, including potassium, to form

NOVEMBER 15, 1949

insoluble soaps which are excreted. Hypokaliemia as well as a calcium deficiency may result.

Prolonged alimentation by intravenous fluids alone may lead to potassium deficiency unless total electrolyte balance is maintained by proper electrolyte composition of the fluids given.

Chronic nephritis usually tends to cause potassium retention. Occasionally, however, potassium loss in the urine is excessive, resulting in potassium deficiency.

## Cracking Joints

EDWARD E. BROWN, M.D.\*

Low temperatures, high humidity, and the ionized air of storms are the principal weather factors predisposing to the cracking sounds heard when joints are flexed or hyperextended.

For four years, Edward E. Brown, M.D., of Ashland, Ore., recorded daily data about the cracking in his phalangeal joints and questioned other subjects. The phenomenon is common in winter, least frequent in summer. Cracking occurs during July in San Francisco, occasionally in Seattle and Portland, and rarely in New York City.

Cracking of joints is probably due to fibrositis. The inflamed fibrous connective tissue contracts; then, when the joints are flexed or hyperextended, the fibers stretch with a cracking noise. The tissue is lengthened after one or more cracks, and no sound is produced until the structures shorten again.

Cracking appears to be associated with nasal obstruction and postnasal drip, conditions also influenced by cool, damp weather.

\* Cause of cracking joints; relation to weather and fibrositis. Northwest Med. 48:537-541, 1949.

INTRAMUSCULAR QUINIDINE GLUCONATE may be used for treatment of cardiac arrhythmia when a rapid effect is desirable or oral administration is not feasible. Samuel Bellet, M.D., and John Urbach, M.D., of the University of Pennsylvania, Philadelphia, find that the preparation is stable and nonirritating and produces typical effects as early as fifteen minutes after injection of 5 to 7.5 gr. Maximum results are obtained in ninety minutes. Quinidine by the intramuscular route does not entail the hazards of intravenous administration and may be lifesaving when quick relief is required from cardiac pain, intense palpitation, dyspnea, vomiting, or shock. I. Lab. & Clim. Med. 34:1118-1120, 1949.

### Endocrine Treatment of Alcoholism

JOHN W. TINTERA, M.D., AND HAROLD W. LOVELL, M.D.\*

St. John's Riverside Hospital Yonkers, N.Y. New York Medical College New York City

ADRENAL cortical extract and a high-fat, low-carbohydrate diet hasten recovery from acute alcoholism. The torturing period of drying-out is practically eliminated and recurrent craving much reduced.

Endocrine treatment was tried by John W. Tintera, M.D., and Harold W. Lovell, M.D., because of the frequent association of hypoadrenocorticism and chronic alcoholism. This adrenal cortex dysfunction seems to be constitutional in some habitual drinkers and acquired in others.

Many young alcoholics are asthenic in physique, with soft, smooth faces and overdeveloped breasts. Adrenal deficiency is shown by hypoglycemia, low 17-ketosteroids, and reduced androgens. Alcoholism is likely to develop at an early age in these subjects, unless strict self-control is exercised.

In a second type of chronic alcoholic, the adrenal cortex and other glands concerned in carbohydrate metabolism have been injured by prolonged intemperance. In most cases of alcoholic cirrhosis the lipoid content of the adrenals is decreased. Endocrine damage may be augmented by worry, grief, or other emotional stimulus.

Though causes differ, the same type of glandular imbalance appears in both types of inebriates. In the acute stage of intoxication lymphocytosis occurs, a further indication of adrenal deficit. During abstinent intervals, hypoglycemia is almost universal. The average fasting blood sugar level is 61 mg. per 100 cc., with a range of 54 to 80 mg. The glucose tolerance curve rises rapidly to moderate elevation, and becomes subnormal in two hours.

Hypoglycemia produces a craving for alcohol and symptoms resembling those of hyperinsulinism. The patient might resort to carbohydrates if he did not know that drinking gives him relief. As more liquor is taken, blood sugar is further reduced, hepatic glycogen is depleted, and fatty infiltration occurs. The liver is no longer

> able to detoxify estrogens, and men undergo feminizing changes.

> Administration of adrenal cortical hormone reverses the entire process. Glycogen is mobilized from tissue protein, blood sugar raised, and hepatic function improved. Psychic and neurologic disorders, epigastric discomfort, and other symptoms subside.



\* Endocrine treatment of alcoholism. Geriatrics 4:274-280, 1949.

For an acute alcoholic state, from three to five days of hospital care is often necessary and always desirable, though out-patient treatment may be successful.

On the first day in the hospital, 30 cc. of adrenal cortical extract is usually given by vein in three doses at intervals of six to eight hours. On the second, 20 cc. is injected in two doses, and for the next three days a single daily dose of 5 to 10 cc. Sedation is seldom required.

After discharge from the hospital, 2 to 5 cc. of the hormone is given intramuscularly twice a week for three weeks, then weekly as long as needed, sometimes for several months.

During and after hospitalization the diet should be high in fat with moderate amounts of protein and little carbohydrate, to prevent sudden fluctuation in the blood sugar level. Vitamin supplements are unnecessary.

Anorexia is overcome by 10 to 15 units of protamine zinc insulin per day injected subcutaneously. For testicular atrophy or other feminizing effects, testosterone propionate may be administered. In women, menopausal symptoms are relieved by testosterone with or without estrogens.

The complete program of Alcoholics Anonymous also should be utilized.

Special psychiatric therapy is reserved for the small group of patients with major psychoses.

RENNIN ESTIMATION of urine may aid in verifying diagnosis of pernicious anemia in patients with gastric achlorhydria without interrupting liver therapy, reports Ole Sylvest, M.D., of Sundby Hospital, Copenhagen. A 10-cc. sample of urine is collected before breakfast, acidified, and compared to a known solution. Rennin content is stated in rennin units per 10 cc. of urine, 1 rennin unit being equal to 0.1 mg. of pepsin Medicinalco (1/3,000). If values above 0.3 rennin units are found, likelihood of pernicious anemia is small.

Acta med. Scandinav. 133:346-349, 1949.

INCIDENCE OF HYPERTENSION in patients with portal cirrhosis is significantly lower than in the general population. These conclusions are reached by Samuel D. Spatt, M.D., and Philip Rosenblatt, M.D., of Jewish Hospital, Brooklyn, after analyzing the blood pressures, heart weights, or both of 60 bed patients over forty years of age who died of portal cirrhosis. Only 18% of the cirrhotic patients had hypertension, although over 50% would be expected to have the disease at that age. Incidence of abnormally heavy hearts was also significantly low among the patients.

Ann. Int. Med. 31:479-483, 1949.

### Radioactive Iodine in Thyroid Diagnosis

D. G. ARNOTT AND ASSOCIATES, LONDON\*

Thyroid dysfunction is shown by radioactive iodine in urine after a tracer dose. Normally, half is concentrated in the gland and the residue excreted.

Healthy, thyrotoxic, and hypothyroid subjects were tested by D. G. Arnott, E. W. Emery, Russell Fraser, M.D., and Q. J. G. Hobson, M.A., of Postgraduate Medical School of London, England.

Potassium iodide, 10 µg. containing 10 microcuries of I<sup>331</sup>, is diluted to 50 cc. As a standard for urine estimation, 5 cc. is pipetted into 20 cc. of 1% potassium iodide solution. The rest of the solution is taken orally after overnight fasting, and 2 rinsings of the container are swallowed.

Urine is collected for forty-eight hours in four samples representing excretion during the first eight hours, the next four, the next twelve, and the final twenty-four hours.

The concentration of radioactive iodine in urine is compared with that in the standard solution by means of the Geiger-Müller counter for liquid samples. Activity in each specimen is expressed as percentage of the dose.

Thyrotoxicosis is shown by low values in the sample taken during the period eight to twelve hours after ingestion. The average level is 0.5%, in contrast to 7% for good health. Hypothyroid determinations for the second day are high, about 20% against a normal 3%.

\* Urinary excretion of radioactive iodine as a diagnostic test in thyroid disease. Lancet 257:460-465, 1949.

HEPATIC INSUFFICIENCY may be ascertained by electrocolorimetric estimation of the turbidity of the patients' serum in distilled water. The method somewhat resembles MacLagan's thymol test, is not so sensitive, and is extremely easy to apply, explain D. Vincent, M.D., of Toulouse, and M. Girard, M.D., of Lyons, France. Four test tubes containing, respectively, 1, 0.5, 0.2, and 0.1 cm. of serum are filled to the top with distilled water. Turbidity is evaluated in one-half hour. The test is positive if the most intense reaction appears in the first two tubes and is above 60 electrocoloric degrees, according to Meunier's scale. The other dilutions do not show flocculation at this time except in very strongly positive cases.

Presse méd. 58:807-808, 1949.

## The Menopausal Syndrome

### ERNEST HOCK, M.D.\*

Charles S. Wilson Memorial Hospital, Johnson City, N.Y.

The vasomotor, neurogenic, and rheumatic symptoms of the menopause may actually be manifestations of urethral causalgia.

Men with prostatic disease have many complaints similar to those of women at the climacteric period. Simple therapeutic measures such as prostatic massage often alleviate the condition almost immediately.

Since the female urethra is the homologue of part of the prostate, menopausal symptoms in women likewise may originate in the urethra, believes Ernest Hock, M.D. An irritative nerve lesion may be caused or aggravated by estrogen deficiency,

which produces degenerative changes in the urethra. Senile urethritis may produce the nerve irritation or may create a susceptibility to a secondary infection. This theory is demonstrated in the diagram.


Organs other than the urethra may contribute to this process. Urethral causalgia cannot explain all the features of the menopausal syndrome. Probably nerve plexuses rather than individual organs should be considered. Hormonal imbalance may possibly in-

crease the reflex irritability of the nervous system, and thereby initiate a causalgia.

Whether the theory is accepted or not, urethral therapy appears to relieve the menopausal discomfort when hormonal treatment will not. Administration of estrogens may have an indirect effect on these symptoms by correcting urethral disease.

Should estrogens and sedatives be ineffective, obvious urinary infection is treated. Otherwise, the amount of relief afforded by passage of a metal sound into the bladder for five to ten minutes should indicate whether the urethra is the source of the symptoms.

Theoretic Mechanism of Menopausal Syndrome



\* The menopausal syndrome. New York State J. Med. 49:2078-2080, 1949.

Cystoscopy may be diagnostic if assay with a sound is inconclusive.

When the urethra and bladder neck are not diseased, 1% novocain solution may be injected into the bladder neck and posterior urethra. Prolonged comfort from this maneuver suggests that the menopausal symptoms in all probability arise in the urethra.

The use of sounds, novocain infiltration, and, occasionally, roentgen irradiation all have therapeutic value in the menopausal syndrome. Patients who have residual urine may be relieved by transurethral resection of the bladder neck.

Many of the symptoms of the male climacteric may also be alleviated by therapeutic treatment of the prostate. This fact suggests that the syndrome is likewise the result of a causalgic process. The rejuvenation which apparently takes place after prostatectomy may also be attributed to the relief of nerve irritation.

### Safe Induction of Pneumothorax

I. J. Selikoff, M.D., I. G. Tchertkoff, M.D., and E. H. Robitzek, M.D.\*

The needle used for initial pneumothorax usually pierces the lung and allows considerable escape of air. To prevent unwarranted laceration, pulmonary tissue should be motionless, the needle small, and insertion no deeper than necessary.

The technic of I. J. Selikoff, M.D., I. G. Tchertkoff, M.D., and E. H. Robitzek, M.D., of the Sea View Hospital, Staten Island, N.Y., is carried out at the bedside. No sedation, cough depressant, treatment room, or special aftercare is required.

The instrument is a 2-cc. or 5-cc. syringe fitted with a sharp, short-bevel needle about 1½ in. long, gauge 22 to 24. The syringe is half filled with sterile saline solution or 2% procaine and the site painted with iodine.

The patient is asked to take and hold a deep breath. The needle is advanced slightly, with constant upward pull on the syringe barrel. If no air bubbles into the solution a few breaths are allowed and the procedure is repeated. When air is obtained, the needle is immediately withdrawn.

In most cases air continues to flow, producing a visible pneumothorax cavity of 150 to 500 cc. Fluoroscopy is done soon after puncture, and six hours later a refill given with a 21-gauge needle.

When no space is seen the induction procedure is repeated; if several attempts fail the pleural space is probably obliterated.

\* Initial pneumothorax: a new safe induction technique. Quart. Bull., Sea View Hosp. 10:93-100, 1949.

### Anatomy and Repair of Varicose Veins

R. STANTON SHERMAN, M.D.\*

University of California, San Francisco

Parlure to eradicate incompetent perforator veins in the legs is the most common cause of persistence or recurrence of varicosities after operation.

Perforator veins in the thigh are much less frequently responsible for poor operative results since, contrary to previous opinion, these veins are seldom incompetent.

After performing numerous anatomic and surgical dissections of the lower limbs, R. Stanton Sherman, M.D., divides the perforating veins of the lower extremity into five main groups (see illustration). These groups

- 1] medial aspect of leg
- 2] lateral aspect of leg
- g] lateral aspect of foot
- 4] short saphenous system
- 5] muscular perforators in the calf

The positions of the perforating vessels are fairly constant. The veins are readily located in terms of distance from the sole of the foot, as indicated in the figures. Some perforating veins pierce only the superficial fascia which extends into the leg from the thigh. Others drain into the deep venous system of the leg by way of intermuscular septa as well as through the deep crural fascia and through the muscles.

Perforator veins are also present on the medial aspect of the foot and ankle but are seldom of clinical significance there.

Optimum therapy of varicose veins should include high ligation of the long saphenous vein, eradication of the mid-Hunter perforator vein in the thigh, and excision of all incompetent vessels in the thigh, the leg, and the foot.

Sclerosing fluid injections may be employed to suppress bleeding. Minor varicosities may be injected locally before surgery.

Careful preoperative examination of the limb is mandatory. Large venous bulbs and palpable fascial defects are noted as frequent sites of varicosities.


Spinal or general analgesia is best employed. After high saphenous ligation, 3 cc. of sclerosing solution is introduced into the saphenous vein. The saphenous vein in the thigh is then explored.

An oblique incision 4 cm. long is made on the inner aspect of the leg just below the knee. The saphenous vein in the leg is stripped digitally and incompetent perforator veins excised. A 3-cm. longitudinal incision is made at the medial border of the tibia about 25 cm. from the sole of the foot. This opening permits an adequate inspection of perforators 3 and 4.

Perforator vein 2 is most common-


<sup>\*</sup> Varicose veins: further findings based on anatomic and surgical dissections. Ann. Surg. 130:218-232, 1949.

### DISTRIBUTION OF PERFORATOR VEINS



Medial aspect of the leg

Lateral aspect of the leg



Posterior leg perforators

Perforator pattern

ly incompetent and is exposed by a separate incision made 19 cm. from the sole and 1 cm. medial to the medial border of the tibia.

Other perforators which are suspected of being incompetent are then investigated by means of separate small incisions when necessary for satisfactory inspection.

Incompetent veins must be exposed extensively to allow the proper deep ligation, otherwise reflux of blood through tributaries will nullify results.

A positive result from a Trendelenburg test-incomplete filling of veins in thirty or more seconds-is inconclusive proof of the absence of incompetent perforator veins. Although in 95% of the cases operated upon the test was positive and competent perforator veins were thereby indicated, in 91% of the legs incompetent perforator veins were found.

## Peptic Ulcer Following Splanchnicectomy

STEPHEN C. MASON, M.D., AND H. M. POLLARD, M.D.\*

Sympathectomy for hypertension interrupts pathways of visceral sensation, including the typical midepigastric pain of peptic ulcer. Chronic ulcers may therefore become worse or new craters develop unnoticed until perforation or massive hemorrhage finally occurs.

Bleeding tendencies may be increased by gastric hyperactivity and by lack of vascular constriction.



Effects of bilateral splanchnicectomy and lower dorsal ganglionectomy were tabulated by Stephen C. Mason, M.D., and H. M. Pollard, M.D., at the University of Michigan, Ann Arbor. Ulcers were present before denervation in 8 patients of a large group and appeared postoperatively for the first time in 4, causing fatal bleeding in 2 of the latter.

If ulcer is active at the time of operation, severe bleeding may occur in a few days. Complications of recent or long-standing ulcers become more frequent and severe after splanchnicectomy.

Vague gastrointestinal complaints should be investigated without delay. An eroding ulcer may be indicated by distress or discomfort, soreness, fullness, sense of pressure, burning, tenderness, dull ache, or nausea and vomiting. Acute pain is usually associated with a serious complication. Immediately before death from perforation the only symptom of one patient was pain in a leg.

\* Peptic ulcer following splanchnicectomy. Surg., Gynec. & Obst. 89:271-284, 1949.

## Physiologic Basis for Burn Treatment

CARL A. MOYER, M.D.\*

Southwestern Medical College, Dallas

TEGLECT of the uninjured parts of a burned or scalded patient may be fatal. Within a matter of minutes, the circulating mass of red blood cells may be reduced as much as 50%, and sodium and salt loss is rapid.

Hemolysis, blood stasis, petechial hemorrhages, and opening and dilatation of capillaries cause a rapid drop

### TREATMENT IN THE THREE STAGES OF THERMAL INJURY

Stage	Physiologic Changes	Treatment
i] Shock and edema for- mation phase Begins with injury and ends 50 to 60 hr. later	Loss of circulating red cells and plasma pro- teins	No scrubbing of burned or scalded parts. This adds to initial injury and aggravates loss of red cells, etc.
		Transfusion of whole blood as soon as possible; give plasma or Hartmann's solution until blood can be obtained; give up to 3 or 4% of body weight in first 24 hr. and continue on subsequent days if pulse is still high and soft
Growing edema of injured parts which induces deficit of extracellular fluid in uninjured parts of body  Danger of ketosis, especially in children	parts which induces defi- cit of extracellular fluid	No water without salt un- less signs of need ap- pear 3 gm, NaCl and 1.5 gm.
	3 gm. NaCl and 1.5 gm. NaHCO <sub>3</sub> per quart of cold water to drink; if vomited while shock ex- ists, give Hartmann's solution intravenously; amounts must be gauged on clinical signs of adequacy or need	
		Food, especially protein. in excess of 100 gm. per 24 hr., and carbohy- drate in excess of 300 gm. per 24 hr.; hard candy

<sup>\*</sup> Recent advances in the chemical supportive therapy of thermal injury. Texas State J. Med. 45:635-639, 1949.

TREATMENT IN THE THREE STAGES OF THERMAL INJURY-(continued)

Stage	Physiologic Changes	Treatment
2] Early recovery phase Starts with subsidence of edema and ends when separation of es- char begins	Delayed hemolysis	Transfusion of blood as required to keep hemo- globin above 10 gm. per 100 cc.
	Edema of injured parts decreasing Tissue growth beginning	Stop salt water by mouth; give water ad libitum Food, including milk and meat, to give 200 to 400 gm. protein daily, and carbohydrate and fat to give 1,000 calories in excess of normal requirements
g] Phase of slough-sepa- ration, granulation, and epithelization	Loss of red cells, especial- ly when dressings are changed	Transfusion of whole blood as for Stage 2
	Rapid expansion of capil- lary vascular bed in granulating areas Sodium loss through gran- ulations	Salt water intermittently as required to make up for loss in excess of sodi- um salts taken in food
		Food as in Stage 2
	Potassium loss through granulations while de- mand by growing cells is great	Never wet dressing with boric acid; it is "poison ous"
	Calcium loss through granulations	Never place patient in water; it also is "poison- ous"
	Protein loss through gran- ulations	Graft as soon as possible

in red blood cells per unit volume of uninjured tissue. If the reduction in the volume of circulating plasma is greater than the decline in volume of circulating red cells per unit of time, the hematocrit reading rises. An initially low hematocrit level indicates that red cells are being lost more rapidly than plasma is.

Extracellular fluid exudes from the undamaged to the burned tissues, first lowering plasma volume and later decreasing extravascular extracellular fluid. Electrolytes in extracellular fluid loss are chiefly sodium chloride and sodium bicarbonate. The reduction

of sodium salts is more serious than the plasma protein decrease.

Supportive measures are essential. Treatment is based on restoration of physiologic changes and is largely directed toward getting whole blood, sodium salts, and water into the patient.

Blood and lactated Ringer's solution (Hartmann's) are immediately infused intravenously to combat shock, urinary suppression, and dehydration. When tolerated, liquids are given orally. A mixture of 3 to 4 gm. of sodium chloride (1 tsp.) and 1.5 to 2 gm. of sodium bicarbonate

or sodium citrate (2/3 tsp.) per quart of water is palatable and averts the acidosis that results from administration of normal saline. Oral saline fluids are discontinued when tissue edema begins to subside (see table).

Water intoxication, points out Carl A. Moyer, M.D., is caused by overloading the severely burned patient with plain water by mouth before adequate amount of salt is given. The water accumulates in the stomach, absorbs sodium salts, and is vomited. The water that held the salts in the plasma remains and dilutes intracellular and extracellular salts. If the fluid is not retained in the stomach but enters the intestine, body salts are still diluted.

Symptoms of headache, tremors, twitching, blurring of vision, vomiting, diarrhea, disorientation, salivation, mania, and even convulsions are caused by excessive water intake and are at times seen in early toxic phases of burns.

No plain water is given by mouth for seventy to one hundred hours unless indicated by severe thirst, dry mucous membranes, rising temperature, and plasma sodium concentration over 138 milliequivalents per liter.

If immersion or massive wet dressings are employed to clean wounds and control infection, the wetting agent must be an isotonic saline solution, 8 gm. of sodium chloride per liter. If water or hypotonic salt solutions are used, sodium salts are drawn into the dressings or bath water, and water enters the body. These changes cause salt-lack shock combined with water intoxication, a rapidly fatal syndrome.

Dressings covering the chest and abdomen should be loose. Pulmonary edema, pulmonary congestion, and atelectasis are easily produced by restriction of breathing. Even a slight impediment to breathing may be fatal to the severely burned.

Ketosis, frequent in children, can be prevented by giving sugar, as hard candy.

Deficiencies of potassium and calcium consequent upon loss from large granulating areas are remedied by high dietary protein, especially milk, meat, and fish.



# Traction and Suspension for Fractures

FRANK E. STINCHFIELD, M.D.\*

Columbia-Presbyterian Medical Center, New York City

The objectives in therapy of compound fractures are reduction and maintenance of alignment, restoration of function of the soft tissues, and protection against additional injury.

In attaining these goals, Frank E. Stinchfield, M.D., finds traction in balanced suspension and fixed traction of great benefit. Traction may be exerted by means of adhesive applied to the skin or by pins or wires through bone.

When skin traction is employed the surface is thoroughly cleansed and the adhesive is smoothed to avoid any wrinkling or exceptation.

An ace bandage is applied in such a fashion that it leaves the upper end of the adhesive exposed so that any slipping may be noted. Skin traction is not recommended for old people, as large areas of tender flabby skin may be avulsed.

When more than a few pounds of traction are required with fractures of the upper extremity, skeletal traction is more comfortable to the patient and more efficient, therapeutically, than skin traction.

For forearm fractures a wire is inserted through the second, third, and fourth metacarpals. The hand is gripped to flatten the metacarpals (Fig. 1a) so that the pin may be inserted as shown in Figure 1b. Figure 1c il-

lustrates incorrect pinning without manual depression of the bones.

The forearm should be in a vertical position with countertraction applied either by a broad sling to the arm with weights attached or by a pin or wire through the upper ulna.

For fractures of the humerus, a wire or pin is inserted through the upper part of the ulna, avoiding the upper cartilaginous area in children. The wire should be inserted from within outward to prevent injury to the ulnar nerve. Traction may be exerted horizontally with the arm parallel to the body, the elbow flexed 90 degrees, and the forearm suspended vertically.

In an alternative method, shown in Figure 1d, the arm is flexed at the shoulder with the forearm over the head and traction is exerted vertically.

For metacarpal and phalangeal fractures, a heavy needle may be passed through the phalanx and countertraction provided by a banjo splint (Fig. 1e, f).

In skeletal traction of the lower limb the following conditions should obtain:

- ► Weights should hang free and cords run easily through different pulleys.
- ➤ Leg and thigh slings should be smooth and always provide equal pressure.
- as shown in Figure 1b. Figure 1c il— The suspension weight should \* The principles of traction and suspension in the treatment of fractures. South. M. J. 42:770-776,

# TRACTION FOR FRACTURES OF UPPER EXTREMITY






FIGURE 1

#### SUSPENSION OF LEG AND




WITH KNEE STRAIGHT



.....

FIGURE 2

#### APPLICATION OF TRACTION



be properly adjusted between the two ends of the half-ring splint, which requires greater force for the upper than for the lower end.

► The weight for knee motion should be within easy reach of the patient, and the trough for the leg should be shallow enough to avoid peroneal pressure.

► The skeletal pin should rest against the bar of the splint instead

of pulling on the skin.

► The inner half of the ring should not press on the groin or perineum.

► The upper support for the splint should be adjusted to allow

proper rotation.

Efficacy of traction and comfort of the patient will be increased if the injured limb is suspended by weights and pulleys different from those used for traction.

Traction applied in balanced suspension eliminates encircling bandages which interfere with blood flow, permits early motion of the adjacent joints, allows use of heat and massage, gives free access to wounds, and lessens the liability of bedsores.

Chief disadvantages of this method of management are that bed rest is usually required and the patient is

not transportable.

The amount of traction required varies with the lapse of time since the injury. With longer time, more force is necessary. Skeletal traction is a very powerful force and excessive traction is too often applied and for too long

Overpull, with separation of fragments, is one of the main causes of delayed union or nonunion. Heavy weight may be applied only until reduction is effected, otherwise results may be unsatisfactory.

Insertion of the pins calls for the same aseptic precautions that are essential in any operation.

The pin that initiates soft tissue movement sooner or later engenders

infection.

A Hodgen or Keller-Blake splint may be used for the lower extremity. The thigh and leg rest in a felt-lined canvas or flannel sling attached to two lateral bars. When a half-ring splint is employed, the ring should be in front and rotated as necessary unless counter pressure against the tuber ischii is deemed advisable (Fig. 20).

The upper margin of the canvas sling should be attached to the ring so that the thigh is always supported. If the band under the fibula is too loose, pressure over the peroneal nerve may cause paralysis.

Figure 2b shows a method of suspension with the knee flexed. The force is applied by adhesive traction on the lower leg, which is supported by a Pearson attachment.

Figure 3a demonstrates how traction weight may be divided between the upper and lower leg by addi-

tion of another weight.

Quite a different system, Russell's method, illustrated in Figure 3b, permits the entire traction and suspension to be accomplished with a single rope, but constant attention is necessary to maintain support of the thigh at all times.

Skeletal traction for the thigh is shown in Figure 3c. The weight controlling knee motion is within reach of the patient and the Pearson at-

tachment is not locked.

### Roseola Infantum

WILLIAM BERENBERG, M.D., CHARLES A. JANEWAY, M.D.

Harvard University, Boston

STANLEY WRIGHT, M.D.\*

University of Rochester, Rochester, N.Y.

NEVER in infancy is one of the most common and difficult problems confronting the general practitioner and pediatrician.

Probably the most frequent febrile exanthem in children is roseola infantum, a disease with a relatively asymptomatic course of three to five days followed by appearance of a morbilliform eruption.

Although William Berenberg, M.D., Stanley Wright, M.D., and Charles A. Janeway, M.D., believe the disease is an almost universal infec-

tion for children under five years of age, few patients are seriously ill or hospitalized. Consequently, medical students do not see many cases and physicians generally are unfamiliar

with the condition.

Fully 95% of the patients are children between six months and three vears of age. During the first months of life the infant may be protected by transplacental passive immunization.

Rarity of roseola infantum after the third year, in view of susceptibility of occasional older patients, raises the possibility that almost all individuals may have the disease before the age of three, probably from contact with adult carriers, and often in an \* Roscola infantum (exanthem subitum). New England J. Med. 241:253-259, 1949.

atypical modified form that defies detection.

An unknown virus is probably responsible. Incubation period is ten to fifteen days, and a single attack confers permanent immunity. Prognosis is excellent, although many sudden deaths of infancy may eventually be proved to be unusually severe, unrecognized forms of roseola.

Complications are rare but can be serious and include convulsions, encephalitis, and hemiplegia.

Infection may develop at any time. Peak incidence is in the spring and fall. Onset of fever is abrupt with the child apparently quite well, active, and alert. In a day or two rose-pink macules appear. The rash does not itch and fades completely without desquamation in two to forty-eight hours.

The temperature is often 103° to 105° when first noted. Fever continues three to nine days. Pulse and respiratory rates increase with temperature. On the first day leukocytosis, by the second or third day, leukopenia, neutropenia, and relative lymphocytosis appear.

Throat and tonsils may be slightly inflamed. Under good light, reddish pinpoint lymphoid elevations are

scen about the uvula and soft palate. Suboccipital and posterior cervical lymph nodes may be palpable by the second or third day of fever and may remain so for several weeks. In 1 of 4 cases, slight catarrhal otitis media develops.

Treatment is purely symptomatic. Sulfonamides and antibiotics are of no value. Rest should be provided and fluids given freely. Aspirin may be prescribed, and if fever is exceptionally high, phenobarbital is administered to prevent convulsions.

## Bronchiectasis in Childhood

C. ELAINE FIELD, M.D., LONDON\*

RREVERSIBLE bronchiectasis can be avoided by careful management of upper respiratory tract disease in childhood.

When bronchial dilatation is established, involved portions of the lung should be removed if enough healthy tissue remains for adequate function. In the experience of C. Elaine Field, M.D., of University College Hospital, London, medical therapy often arrests but seldom reverses the progress of disease. Surgical treatment, on the other hand, may be curative in half the cases.

Results of treatment were observed in 202 children for periods up to ten years. Bilateral lobectomy was successful in 10 instances. Complete resection of affected tissue was curative in 54% of cases; medical treatment in 12%.

During recovery from pneumonia, pertussis, and other respiratory infections the chest should be examined repeatedly. Asthma with bronchitis and so-called chronic bronchitis may actually comprise early bronchiectasis. If a cough persists, postural drainage and breathing exercises should be employed.

When constant cough and dilatation have continued for a year or more, surgical treatment may be necessary. In extensive cases, grossly diseased parts may be excised and symptoms relieved by other measures.

Regardless of slightness or severity of manifestations, operation should be done in all cases not too advanced for complete anatomic and symptomatic cure. If possible, segments are removed rather than entire lobes, and half the pulmonary tissue is saved.

In a child, the remaining bronchi grow and lungs occupy almost the same area as before. Much more function returns to children after operation than to adults.

The most effective chemotherapy combines penicillin with streptomycin.

\* Bronchiectasis in childhood. Pediatrics 4:355-372, 1949.

# Tonsils, Adenoids, and Allergy

NORMAN W. CLEIN, M.D.\*

University of Washington, Seattle

REGROWTH of adenoid and tonsillar tissue after resection is usually due to unsuspected allergy.

Before tonsillectomy and adenoidectomy are undertaken for nasopharyngeal disorders, an allergic source should be sought.

Norman W. Clein, M.D., believes that operation may fail in presence of any of the following symptoms:

- ► Constantly running nose more troublesome at night and on arising
- ► Repeated colds and chronic cough, also worse at night
- ➤ So-called chronic sinusitis with or without migraine
- ► Persistence of complaints after operation and return of hypertrophic tissue.

In about half the allergic cases, pathologic tissue shrinks or atrophies during treatment and no surgery is required. In the remainder, surgery may be done, since postoperative hypertrophy will be negligible and return of the original symptoms less likely

Nasopharyngeal allergy is manifest in children by symptoms of the following two types:

1 One cold after another, usually without fever, seasonal or perennial. Mouth kept open most of the time, especially at night. Habitual snoring. Sniffling and nose blowing, notably early in the morning.

\* Allergy and the tonsil problem in children. Ann. Allergy 7:329-333, 1949.

2 Colds accompanied by hacking or clearing of the throat. Deep, hard, dry cough, aggravated by exercise or fatigue and more frequent at night. Wheezing and difficult breathing. Attacks of croup, sometimes accompanied by respiratory infection. Lowgrade fever occasionally continuing for months.

An excess of eosinophils in repeated nasal smears is pathognomonic of the allergic cold.

After unsuccessful operation, lymphoid tissue returns more often to the nose than to the throat, and the most common complaint is mouth breathing. Anterior turbinates may be swollen or normal, but invisible postnasal edema is often shown by blowing on one side. Unlike stenosis from a large adenoid mass, allergic congestion subsides at intervals.

Tonsillar hyperplasia may appear as early as five months after resection. Reddish swollen lymphoid areas resembling granulation tissue may be observed under the anterior pillar or in the middle of a fossa. Lateral pharyngeal walls often contain scattered, round, raised nodules of pinhead to pea size.

Operation during the pollen season may aggravate hay fever or initiate asthma.

Postnasal lymphoid hyperplasia may be reduced by application of radium.

# Development of Vision in Childhood

ARNOLD GESELL, M.D.\*

Yale University, New Haven, Conn.

Like speech, vision develops slowly and passes through many contrasting stages. Since all are closely related to posture, manual skills, motor habits, intelligence, and personality, one atypical trait often points to others.

Arnold Gesell, M.D., advises systematic preschool examinations.

Brain lesions as well as ocular defects may be revealed and future problems forecast. At all ages both educational and corrective measures should be geared to the child's actual capacity.

The relation of sight to behavior was investigated from premature

birth to the tenth year in the home, guidance nursery, and school. About 50 children were observed at each of twenty age levels.

When the newborn baby sees something of interest, body activity tends to subside. A full-term infant begins to notice a near-by, approaching object on the first day of life, sustains fixation in the first week, and regards more distant objects in a month. He will notice a pellet 7 mm. in diameter twenty weeks sooner than he can pick it up with his fingers.

Seeing ability is shown by ocular and other attitudes. Excessive hand

regard beyond the age of twelve weeks may signify myopia, while delayed hand regard may indicate mental deficiency or atypical spatial manipulation. Unusual eye-hand patterns are sometimes harbingers of speech defects and reading disability. A clumsy hand or foot at the age of two and a half years may foretell strabismus a

year or two later.

The preschool child's use of cup, spoon, crayon, paints, and picture books is significant. Too much or too little staring, caution, or withdrawal from companions may be due to visual rather than emotional difficulty.

In the school years, changing function causes many problems. Sustained attention may be demanded of a pupil in the phase of shifting activity, or ability rated poor because instruction is too advanced. Premature reading and writing are to be avoided. Frequently, inadequate vision requires improved teaching rather than treatment.

Sight operates in three basic fields: skeletal, visceral, and cortical. The skeletal component seeks and holds a visual image, the visceral component discriminates and defines, while cortical activity unifies and interprets.

All three functions vary with ad-

The developmental aspect of child vision. J. Pediat. 35:310-316, 1949.



vancing maturity. After the age of five years new abilities appear, not always in balanced or well-timed relation. At seven years the visceral-skeletal linkage is relatively tight, at eight more facile, and at nine years again consolidated.

Visual ability also depends on coordination of the two eyes and right versus left dominance; reach to different planes of regard; scope of central and peripheral view; and drift, including span of ocular ductions, preferred zones of regard, accommodation, and dominant direction.

The retinoscope shows a close relation between visual function and total activity. As the eye gropes, grasps, and manipulates, the reflected beam of light varies in brightness, motion, direction, speed, and color. Light intensifies when the infant seizes a target.

Not only refraction and fusion but the child's organization of visual equipment should be considered in overcoming disability. Partial correction may call forth the child's best resources when full correction would supply a hampering crutch. Training and therapeutic regimens should be wisely timed in short, interesting sessions and perfectionist aims should be avoided.

# Histamine Therapy with Electric Shock

A. M. SACKLER, M.D., M. D. SACKLER, M.D., AND R. R. SACKLER, M.D.\*

PRELIMINARY course of histamine injections may double the number of schizophrenic, manic depressive, and involutional psychotic patients leaving mental hospitals after electric convulsive therapy. A. M. Sackler, M.D., M. D. Sackler, M.D., and R. R. Sackler, M.D., of the Creedmoor State Hospital, New York City, inject histamine base in saline solution subcutaneously, intramuscularly, or intravenously in doses of 1 to 3 mg. once or twice daily for a few days to several weeks.

A group of 25 women refractory to these injections received electrotherapy. Nearly half the subjects improved and 16% were able to go home. Of those receiving histamine or electric convulsive therapy alone, about one-fourth were benefited and one-eighth discharged.

Histamine is most effective when administered early. The majority of women improved by drug or combined treatment had been in the hospital not more than six months. Of the short-term group, 31% were able to return-home after combined therapy, 21% after histamine only, and 14% after shock.

\* Nonconvulsive biochemotherapy with histamine and electric convulsive therapy, J. Nerv. & Ment. Dis. 110:185-197, 1949.

# Office Treatment of Syphilis

THEODORE ROSENTHAL, M.D.\*

Columbia University and Department of Health, New York City

In private practice the number of patients treated for syphilis is relatively small. Trial-and-error schedules used in large clinics are not applicable since the practitioner desires to give sufficient medication to insure the best possible results for each individual.

Theodore Rosenthal, M.D., outlines for the various forms of syphilis the following treatment, which, in the light of present knowledge, approximates minimal therapy but assures optimum results. Although penicillin is the principal drug employed, the addition of arsenoxide and bismuth has genuine synergistic value.

Primary and secondary syphilis and latent and benign late syphilis are best treated as follows: 6,000,000 units of procaine penicillin G is administered intramuscularly in doses of 600,000 units each day for ten days except Sundays. Then twenty injections of mapharsen are given at the rate of two injections per week for ten weeks. One injection of bismuth is given each week for ten weeks either concurrent with or following the course of mapharsen.

Thereafter, titrated serologic tests for syphilis must be done monthly for one year and patients carefully observed.

Spinal fluid examination is performed before therapy in late syphilis eight to fifteen days.

\* Treatment of syphilis in private practice. New York Med. 5:16-18, 34, 37, 1949.

and within twelve months after treatment in either primary or secondary syphilis.

The entire course is repeated for either a clinical or serologic therapeutic failure.

If a patient with neurosyphilis can be ambulatory, 9,000,000 units of procaine penicillin G in doses of 600,000 units intramuscularly every other day for fifteen injections is the treatment suggested.

Consultation with a neurologist is desirable for patients with neurosyphilis requiring hospitalization.

Cardiovascular syphilis must be treated cautiously, beginning with heavy metals for six to eight weeks. At least 6,000,000 units of penicillin in relatively small divided doses is then given over a period of fifteen or more days.

Therapy for ocular syphilis should always be planned with an ophthalmologist.

A serologic pattern may not be established for ten to twelve weeks in the newborn. For congenital syphilis penicillin G is administered in aqueous solution if the patient is less than two years of age.

A total amount of from 100,000 to 400,000 units per kilogram of body weight is given in divided doses throughout the day and night for eight to fifteen days. Older children may receive procaine penicillin G in doses of 6,000 to 12,000 units per kilogram of body weight daily for two to three weeks.

Syphilis is almost certainly prevented in the offspring if the syphilitic pregnant woman receives adequate

penicillin therapy.

A pregnant woman with syphilis is given 600,000 units of procaine penicillin G daily for ten days, beginning immediately after the diagnosis is made, regardless of the stage of gestation, even as late as when admitted to the hospital for delivery. Monthly

titers are determined until delivery. The patient probably should be given penicillin during each successive pregnancy even when the infection is apparently cured.

Penicillin-intolerant patients require prolonged courses of mapharsen and bismuth, with a minimum of forty injections of mapharsen and sixty of bismuth. Urine should be examined periodically and attention given to optic or auditory nerve involvement, severe headaches, dermatitis, jaundice, enlargement of the liver, or acholic stools.

SARCOIDOSIS apparently is not significantly affected by vitamin D<sub>2</sub> therapy. Remissions appear in less than a year with or without treatment in 50% of cases, finds Carl T. Nelson, M.D., of Columbia University, New York City, after a twelve-month study of 16 patients with the disease, 8 of whom were given large doses of calciferol and 8 of whom were untreated. Although recovery was more rapid in the patients given calciferol, the difference was not great. Vitamin D<sub>2</sub> dosage was 120,000 units daily in a mixture of equal parts of ethyl alcohol and propylene glycol unless improvement was noted, in which event the dosage was halved. Previous reports on the efficacy of calciferol for sarcoidosis have concerned white patients. The fact that 15 of the 16 patients were Negroes may account for the disparity in results.

J. Invest. Dermat. 13:81-84, 1949.

DISSEMINATED NEURODERMATITIS may be alleviated by topical application of Thephorin. The antihistaminic drug apparently is able to penetrate skin rapidly, since relief from itching usually starts within fifteen minutes, find Wilfred E. Wooldridge, M.D., and Herbert L. Joseph, M.D. Effects last two to three hours or longer. Nearly all of 23 patients in the Barnard Free Skin and Cancer Hospital and Washington University, St. Louis, were subjectively relieved by the ointment although the eruptions of only 2 subjects were completely cleared. A single case of induced sensitivity occurred, but no constitutional reactions were noted. Results were less successful with circumscribed neurodermatitis.

Arch. Dermat. & Syph. 60:390-403, 1949.

# The Eye in General Practice

A. J. BALLANTYNE, M.D.\*

University of Glasgow, Scotland

The family physician is often confronted with ocular problems such as sudden loss of sight, uveitis from infectious disease, and the retinal vascular changes of essential hypertension, diabetes, and pregnancy.

A. J. Ballantyne, M.D., lists a number of helpful diagnostic and prognostic clues.

#### LOSS OF SIGHT

Few symptoms cover a larger field of possible causes than sudden failure of vision.

Detachment of the retina is frequently responsible when sight of one eye is lost. At whatever point the retina separates, fluid tends to settle in the lower part of the eye, throwing a dark curtain or shadow over the upper part of the visual field. Test finger movements may show the defect before ophthalmoscopic examination is made.

Sudden darkening of the whole field of vision in an eye most often results from hemorrhage into the vitreous, ranging in severity from a reddish tinge to obliteration of light. Repeated minor hemorrhages with intervals of apparent recovery are sometimes caused by tuberculous changes. In many cases blood clots organize, the retina is detached, and the eye is lost.

Complete obstruction of a central retinal vessel causes practically total blindness. With arterial embolism the entire retina rapidly becomes ischemic and, since viability is destroyed in a few minutes, rarely regains function. Occasionally the clot is moved or absorbed.

Venous obstruction is productive of edema with multiple large hemorrhages throughout the retina and usually abolishes useful sight.

Amblyopia from arterial spasm tends to be sudden, short, recurrent, and sooner or later bilateral. With arteriosclerosis and hypertension, a fatal cerebral seizure may ensue.

Retrobulbar optic neuritis causes a central scotoma often preceded or followed by transient diplopia. In some cases eye movements and pressure on the eyeball are painful. Recovery is the rule.

Sudden binocular loss of sight is ordinarily due to a toxin. Amblyopia or amaurosis during pregnancy may be a serious forewarning of eclampsia or, less often, a consequence of hyperemesis gravidarum with retrobulbar neuritis. In the latter case, large doses of vitamins B<sub>1</sub> and C may be helpful.

Sudden temporary dimness of vision with early diabetes is due to change in the crystalline lens. The eyes become myopic but, with system-

<sup>\*</sup> The eye in general medicine. Practitioner 163:173-181, 1949.

ic treatment, revert to the original refractive state.

Quinine blindness is rare and apparently due to idiosyncrasy. Wide inactive pupils, deafness, and impaired consciousness may be observed. Although the retina becomes ischemic, central and incomplete peripheral vision returns.

#### INFLAMMATORY OCULAR LESIONS

Iritis, cyclitis, and choroiditis may occur with the exanthems, focal sepsis, syphilis, and other systemic diseases. Subacute uveitis may be the outstanding manifestation of Boeck's sarcoidosis or of spondylarthritis ankylopoietica. When the cause of uveitis is not found, antituberculous measures are often effective.

Congenital cataract and related deformities may result from maternal infection with mumps, measles, chickenpox, and scarlet fever as well as rubella in early pregnancy.

#### HYPERTENSION

In the first stage of essential hypertension, retinal vessels are well filled and tortuous, with good color and copper high lights. Veins are slightly compressed at arterial crossings. Later the arteries narrow and the veins become more obviously dented.

Superficial striate or spindle-shaped hemorrhages occur about the disk and eventually bleeding involves all retinal layers. Exudates appear as small discrete points which coalesce, scattered woolly white patches, or diffuse serofibrinous infiltration. A macular star is seen in the fully developed stage. Arteriosclerosis may develop before or after onset of malignant hypertension but is usually prominent in the final stage.

In most cases the prognosis of hypertension corresponds to degree of eyeground change. Early retinal lesions without hemorrhages or exudates may disappear with treatment and occasionally severe changes are reversed by sympathectomy.

#### DIABETES

Diabetes mellitus first produces capillary aneurysms seen as minute red dots near the macula. Round or polygonal hemorrhages then appear in deep retinal layers, and punctate exudates form increasingly larger patches with a waxy yellowish surface.

Retinal veins become full and tortuous, with kinks, loops, and fatty infiltration of walls. Networks of new vessels develop. Retinopathy is not related to severity of diabetes. Eyeground lesions steadily progress and may cause blindness in a reasonably healthy patient.

#### PREGNANCY

Ophthalmoscopic examination during pregnancy may show vessel changes of incipient hypertension, arteriosclerosis, nephrosclerosis, or diabetes. Early fundus changes of hyperemesis gravidarum consist of hemorrhages and papilledema.

Preeclamptic toxemia with profound visual loss may produce only diffuse narrowing of retinal arteries and arteriovenous crossings with slight edema of the disk and retina. If fundal lesions do not improve after two weeks of general therapy, pregnancy should be terminated at once.

# The Eye in General Practice

A. J. BALLANTYNE, M.D.\*

University of Glasgow, Scotland

The family physician is often confronted with ocular problems such as sudden loss of sight, uveitis from infectious disease, and the retinal vascular changes of essential hypertension, diabetes, and pregnancy.

A. J. Ballantyne, M.D., lists a number of helpful diagnostic and prognostic clues.

#### LOSS OF SIGHT

Few symptoms cover a larger field of possible causes than sudden failure of vision.

Detachment of the retina is frequently responsible when sight of one eye is lost. At whatever point the retina separates, fluid tends to settle in the lower part of the eye, throwing a dark curtain or shadow over the upper part of the visual field. Test finger movements may show the defect before ophthalmoscopic examination is made.

Sudden darkening of the whole field of vision in an eye most often results from hemorrhage into the vitreous, ranging in severity from a reddish tinge to obliteration of light. Repeated minor hemorrhages with intervals of apparent recovery are sometimes caused by tuberculous changes. In many cases blood clots organize, the retina is detached, and the eye is lost.

Complete obstruction of a central retinal vessel causes practically total blindness. With arterial embolism the entire retina rapidly becomes ischemic and, since viability is destroyed in a few minutes, rarely regains function. Occasionally the clot is moved or absorbed.

Venous obstruction is productive of cdema with multiple large hemorrhages throughout the retina and usually abolishes useful sight.

Amblyopia from arterial spasm tends to be sudden, short, recurrent, and sooner or later bilateral. With arteriosclerosis and hypertension, a fatal cerebral seizure may ensue.

Retrobulbar optic neuritis causes a central scotoma often preceded or followed by transient diplopia. In some cases eye movements and pressure on the eyeball are painful. Recovery is the rule.

Sudden binocular loss of sight is ordinarily due to a toxin. Amblyopia or amaurosis during pregnancy may be a serious forewarning of eclampsia or, less often, a consequence of hyperemesis gravidarum with retrobulbar neuritis. In the latter case, large doses of vitamins B<sub>1</sub> and C may be helpful.

Sudden temporary dimness of vision with early diabetes is due to change in the crystalline lens. The eyes become myopic but, with system-

<sup>\*</sup> The eye in general medicine. Practitioner 163:173-181, 1949.

ic treatment, revert to the original refractive state.

Quinine blindness is rare and apparently due to idiosyncrasy. Wide inactive pupils, deafness, and impaired consciousness may be observed. Although the retina becomes ischemic, central and incomplete peripheral vision returns.

#### INFLAMMATORY OCULAR LESIONS

Iritis, cyclitis, and choroiditis may occur with the exanthems, focal sepsis, syphilis, and other systemic diseases. Subacute uveitis may be the outstanding manifestation of Boeck's sarcoidosis or of spondylarthritis ankylopoietica. When the cause of uveitis is not found, antituberculous measures are often effective.

Congenital cataract and related deformities may result from maternal infection with mumps, measles, chickenpox, and scarlet fever as well as rubella in early pregnancy.

#### HYPERTENSION

In the first stage of essential hypertension, retinal vessels are well filled and tortuous, with good color and copper high lights. Veins are slightly compressed at arterial crossings. Later the arteries narrow and the veins become more obviously dented.

Superficial striate or spindle-shaped hemorrhages occur about the disk and eventually bleeding involves all retinal layers. Exudates appear as small discrete points which coalesce, scattered woolly white patches, or diffuse serofibrinous infiltration. A macular star is seen in the fully developed stage. Arteriosclerosis may develop before or after onset of malignant hypertension but is usually prominent in the final stage.

In most cases the prognosis of hypertension corresponds to degree of eyeground change. Early retinal lesions without hemorrhages or exudates may disappear with treatment and occasionally severe changes are reversed by sympathectomy.

#### DIABETES

Diabetes mellitus first produces capillary aneurysms seen as minute red dots near the macula. Round or polygonal hemorrhages then appear in deep retinal layers, and punctate exudates form increasingly larger patches with a waxy yellowish surface.

Retinal veins become full and tortuous, with kinks, loops, and fatty infiltration of walls. Networks of new vessels develop. Retinopathy is not related to severity of diabetes. Eyeground lesions steadily progress and may cause blindness in a reasonably healthy patient.

#### PREGNANCY

Ophthalmoscopic examination during pregnancy may show vessel changes of incipient hypertension, arteriosclerosis, nephrosclerosis, or diabetes. Early fundus changes of hyperemesis gravidarum consist of hemorrhages and papilledema.

Preeclamptic toxemia with profound visual loss may produce only diffuse narrowing of retinal arteries and arteriovenous crossings with slight edema of the disk and retina. If fundal lesions do not improve after two weeks of general therapy, pregnancy should be terminated at once.

# Surgery for Primary Glaucoma

CONRAD BERENS, M.D., L. BENJAMIN SHEPPARD, M.D., ARTHUR B. DUEL, JR., M.D., AND LOUIS J. GIRARD, M.D.\*

New York City

SELECTION of the best operation for primary glaucoma is difficult because the disease has no universally recognized classification. The principal types of glaucoma, narrow and wide

angle, should be differentiated, but even after careful study the two are not always distinguishable.

Conrad Berens, M.D., L. Benjamin Sheppard, M.D., Arthur B. Duel, Jr., M.D., and Louis J. Girard, M.D., observe, however, that different technics have been successful in treating glaucoma and that choice of treatment continues to be empiric, depending much on the surgeon's preference and ability.

Aspects of each case, including base pressure, angle width, peripheral anterior synechiae, chronic infection, and atrophy of the iris and conjunctiva, must be taken into consideration before choice of operation is made.

For acute primary glaucoma, operation is ordinarily indicated if medical therapy is ineffective after twelve hours' trial. Delay allows peripheral anterior synechiae to form.

The best technic is usually a basal iridectomy, especially when surgery is performed early and the angle is narrow. The same procedure may also been used successful to the surgery of primary glaucoma. South. M. J. 42:731-738, 1949.

be advisable with chronic primary glaucoma if the angle is narrow, the iris not atrophic, base pressure low, and few peripheral anterior synechiae are evident. Even when some of these conditions

are unfavorable, the operation may succeed in skilled hands.

Basal iridectomy, even when combined with early and repeated massage, sometimes fails to control tension, especially when the anterior chamber is deep, previous medical therapy has not reduced tension, and anterior synechiae are seen or suspected. In such cases iridocorneosclerectomy may be done.

For wide-angle glaucoma, an external or internal filtering operation should be chosen. Cyclodialysis with the introduction of air is used if the base pressure is low and the fields strongly contrasted. When pressure is over 40 mm. Hg (Schiötz), iridocorneosclerectomy is advisable; the iris pillars are incarcerated in the wound if pressure is above 50 mm.

With a rapidly developing cataract, complete iridectomy is performed.

Although iridencleisis is more effective than iridectomy in the later stages of acute primary glaucoma and has been used successfully in severe chronic cases, the operation may provoke sympathetic ophthalmitis.

Not enough is known about cycloelectrolysis to recommend the operation as a primary procedure, but the technic may be used when other operations have failed to control tension and is the best procedure for Negro patients, in whom filtering operations frequently fail. If tension is high after operation in these cases, aspiration or electroparacentesis is used.


Gun-barrel fields are probably a contraindication to ophthalmologic operation.

# Sterile Urinary Drainage

WESTON T. BUDDINGTON, M.D., AND ROGER C. GRAVES, M.D.\*

M ECHANICAL introduction of bacteria is the usual cause for urinary infection in patients subjected to prolonged catheter drainage. Cystitis from urethral or suprapubic procedures may be much reduced or completely averted by consistently

sterile technic.



Weston T. Buddington, M.D., and Roger C. Graves, M.D., of Tufts College, Boston, prevent reflux of contaminated urine with a special drainage bottle (see illustration).

The stopper has two holes for a connecting tube and an air vent. Bottle, cork, rubber tubing, and glass connections are autoclaved

as a unit and replaced every other day.

The sterile drainage system has two advantages, both of which prevent retrograde extension of bacteria along the catheter:

1] When the container is raised above bladder level, as often done by nurse and patient, reverse siphonage of bottle contents is practically impossible.

2] If backflow does occur, the pool of urine is sterile and does not contaminate the lower end of the tubing.

In obtaining specimens for culture, a section of drainage tube is prepared with tincture of iodine, and urine is aspirated with needle and syringe.

Antibiotic treatment is continued night and day with sulfadiazine, penicillin, methenamine with sodium acid phosphate, or other drugs. Streptomycin is given for severe infection with gram-negative bacilli, especially *Proteus* and *Pseudomonas*, and just before discharge from the hospital.

\* Management of catheter drainage. J. Urol. 62:387-393, 1949.

# Cause and Therapy of Pelvic Myalgia

NORBORNE B. POWELL, M.D.\*

Baylor University, Houston

Painful spasm of the piriformis muscle group may arise from posterior urethritis or from anorectal and orthopedic disorders. As a result, bizarre unilateral or bilateral symptoms may appear in one or more regions of the pelvis. hip, or thigh.

The principal causes of pelvic myalgia are chronic prostatitis in males and the urethral syndrome in females. Norborne B. Powell, M.D., diagnoses the condition by rectal or vaginal palpation. Tense muscles can usually be relaxed by massage of the pelvic fioor.

Chronic inflammation of the urethra, prostate, or rectum is accompanied by lymph stasis along the posterior pelvic floor, which produces cdema of the common peroneal nerve.

The piriformis muscle, which is usually pierced by the nerve, may then tighten, compressing the sciatic nerve between the piriformis and superior gemellus. Consequent slight pressure on sensory fibers results in pelvic myalgia. Pain from skeletal or postural defects is probably due to mechanical stretching of muscles.

The piriformis, inferior and superior gemelli, obturator internus, gluteus medius, levator ani, and coccygeus may be affected singly or in groups. Tension produces symptoms

m areas supplied by the fourth and fifth lumbar nerves and the first and second sacral nerves.

Pain, ache, or numbness may occur about the hip joint, coccyx, or both; indefinite ache down the sciatic area; discomfort or ache in one or both groins; vague distress or sense of weight in the vagina, rectum or deep pelvis of a woman, in the perineum or rectum of a man.

When a finger is inserted into the rectum or vagina a tender muscular ridge is felt from the coccyx to the medial surface of the hip joint. Spasm confined to one side is particularly obvious.

Although urethral therapy would eventually relieve the spastic condition, cure is hastened by gentle massage. The patient should be warned that in 10% of cases symptoms temporarily become worse during the course of treatment.

The finger is placed in vagina or rectum and swept laterally from coccyx to acetabulum and back several times. When the entire surface of the muscular ridge has been covered, direction is changed. Stroking continues in a plane perpendicular to the first maneuver, gradually working toward the acetabulum.

Massage seldom produces more than slight discomfort, but if tension is extreme the lightest touch may

\* Pelvic myalgia: a complication of posterior urethritis in males and females. J. Urol. 62:245-249.

cause excruciating pain. After a momentary pause, however, treatment can generally be resumed.

As a rule, myalgia partly or completely disappears for two to four days after the first treatment. Exacerbations are checked by aspirin, codeine, and external heat. The massage should be repeated at intervals until symptoms are gone.

# Bell's Palsy

JEROME A. HILGER, M.D.\*

 $\Gamma$  or want of an etiologic description, peripheral facial paralysis not caused by trauma, infection, or neoplasm retains the designation of Bell's palsy.

The condition is an ischemic neuritis which results from segmental arteriolar spasm, observes Jerome A. Hilger, M.D., of St. Paul. Edema of the nerve may occur as a secondary phenomenon. The tendency to arteriolar constriction is probably on the basis of an inherited potential imbalance of the autonomic nervous system.

Unusual chilling of the side of the face has been cited as a precipitating cause of the paralysis in many cases. Severe emotional upset or shock may be the precipitating factor. A familial tendency exists.

Onset is usually sudden. Pain about the ear is not uncommon in the first forty-eight hours. Adults are the most frequently afflicted. The paralysis may be partial or complete. The only prognostic sign is the completeness of paralysis or the loss of faradism. Hyperacusis and loss of taste are common in early stages, but these sensory disturbances usually disappear even if the paralysis persists. This is due to collateral arterial supply to the proximal portion of the nerve trunk.

Principal fault in therapy today is the neglect of nonsurgical measures. Recovery occurs spontaneously in about 85% of cases but can be greatly facilitated by active therapeutic measures.

Therapy should be directed toward relief of the vasospasm. Immediate use of vasodilators, such as nicotinic acid, intravenous histamine, or papaverine, or sympathetic paralyzers, such as ergotamine tartrate or tetraethylammonium chloride, may be effective. Cervical sympathetic block is sometimes valuable.

Surgical decompression of the edematous nerve trunk may be necessary in some cases, but the timing of the procedure has not been definitely determined. Waiting until fibrosis replaces necrosis in the nerve trunk is certainly unwise; however, fibrosis is infrequent in nerves with paresis of less than six months' duration. Muscle reactivity may be retained through electrical stimulation, support, and massage.

\* The nature of Bell's palsy. Laryngoscope 59:228-235, 1949.

# Cerebral Apoplexy

HARRY M. ZIMMERMAN, M.D.\*

Columbia University, New York City

DEATHS from brain hemorrhage are often prevented and disabling sequelae reduced by evacuation of blood. Before exploratory craniotomy is undertaken, however, cerebral hemorrhage must be differentiated from

infarct, the lesion accurately located, and pathogenesis understood.

Both types of cerebrovascular accident were investigated by Harry M. Zimmerman, M.D. Brains were examined post mortem in 182 consecutive cases; of these, 107 revealed spontaneous hemorrhages, and 75 had infarcts. Effects of 175 nonfatal cerebrovascular lesions were also reviewed; here the two conditions were about evenly divided.

Brain hemorrhage occurs with equal frequency in men and women, and incidence of fatal and nonfatal lesions reaches a peak in the sixth decade of life. Most deaths occur during the first attack, within five weeks of onset.

The cause of bleeding is practically always hypertension, with heart and kidney involvement in the majority of instances.

The symptoms of hemorrhagic stroke, in order of frequency, are sudden loss of consciousness, paralysis,



a fall, vomiting, headache, vertigo, convulsions, and weakness.

The delayed effects are unconsciousness, reflex changes, paralysis, pupillary or fundic abnormalities, a positive Babinski sign, spasticity, and vomiting.

As a rule, spinal fluid contains blood, and pressure is increased.

Infarction of the brain is 2 or 3 times more common in men than in women. Peak incidence is at least ten years later than with brain hemorrhage. Prospects of survival are better than with bleeding; fewer fatalities occur in the first attack and a larger number three years or more after onset of symptoms.

Hypertension is relatively infrequent, developing before death in less than half the cases, often without cardiovascular renal disease.

Infarction is first shown by a generalized paralysis, unconsciousness, mental disturbance, a fall, weakness, or dysarthria. Most common during the subsequent illness are paralysis, unconsciousness, reflex changes, mental aberration, dysarthria, pupillary disorders, a positive Babinski sign, and nuchal rigidity.

Spinal fluid seldom contains blood, and pressure is generally normal.

 \* Cerebral apoplexy: mechanism and differential diagnosis. New York State J. Med. 49:2153-2157, 1949. Spontaneous cerebral hemorrhage occurs in right and left cerebral hemispheres with equal frequency. The site in 38% of cases is supplied by the lenticulostriate artery or branches. Apart from basal ganglia, affected regions may include the frontal lobes, cerebellum, calcarine areas, ventricular system, occipital poles, pons, and occasionally the centrum ovale of all lobes in one hemisphere.

About three-fourths of hemorrhages are due to rupture of miliary dissecting aneurysms in atherosclerotic cerebral arterioles. From intramural to massive flow is but a step. As blood spreads into nervous tissue, neighboring capillaries and venules are torn and add other hemorrhages. The tide may reach Virchow-Robin spaces of

larger vessels and mix with cerebrospinal fluid or burst into the subarachnoid region through disrupted gray matter. The ventricles may be flooded by a break in the basal ganglia.

The hematoma apparently reaches maximum size in minutes or at most a few hours, then forms a permanent cavity by destruction or compression of adjacent zones. After remaining liquid for several days, blood clots to the consistency of soft currant jelly. Venous stasis, distant parenchymal injury, and herniation of the cerebellar tonsils may result. If death is escaped, the swelling subsides in time and blood is absorbed, but the cavity remains. Even in the course of years, the glia makes little attempt at repair.

SIGMOID VOLVULUS can often be identified by a unique radiologic sign. If the ileocecal valve is incompetent, the two parts of the obstructed sigmoid coil are uniformly expanded to the same diameter, observes M. Arias Bellini, M.D., of Montevideo, Uruguay. With a competent valve the colon is also blocked and expanded, but the cecum and ascending colon are much wider than the descending portion. Roentgenograms in standing and supine positions sometimes suffice for diagnosis. A barium enema may indicate sigmoid volvulus if the barium remains in the rectosigmoid region or passes the obstruction and enters the distended coil, from which the medium cannot be expelled.

Radiology 53:268-270, 1949.

BRONCHIAL STONES are more common than heretofore suspected. Using laminagraphy, Eugene Freedman, M.D., and James H. Billings, M.D., found 7 instances in one year at Cedars of Lebanon Hospital, Los Angeles. The usual source is perforation of calcified tuberculous lymph nodes. Concretions should be sought when paroxysmal cough is associated with bronchial obstruction, pulmonary suppuration, or hemorrhage if radiography shows calcific shadows in consolidated areas.

Radiology 53:203-215, 1949.

# Medical Forum

Discussion of articles published in Modern Medicine is always welcome. Address all communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.

#### Gastric Suction after Cesarean Delivery\*

TO THE EDITORS: I believe that gastric suction of infants should be routine after cesarean section, as do Drs. Sydney S. Gellis, Priscilla White, and William Pfeffer. I have used it recently and have been surprised at the amount of fluid obtained.

CECIL W. SEIBERT, M.D.

Waterloo, Iowa

TO THE EDITORS: I do not think gastric suction on babies born by cesarean section should be done routinely, as suggested by Drs. Sydney S. Gellis, Priscilla White, and William Pfeffer. This procedure is unnecessary in a great majority of cases, and is actually traumatic in many.

Obstetricians are impressed with the high incidence of respiratory difficulty, asphyxia, and strangulation in these babies as compared with those delivered by the vaginal route.

Several months ago, I heard Dr. Fred H. Falls of Chicago express the opinion that as the baby passes along the birth canal and over the perineum in vertex presentation, the chest is squeezed in such a manner as to force from the bronchial tree, and to some \*Modern Medicine, July 15, 1949, p. 64.

extent the other air passages, mucus and amniotic fluid that has collected. This does not occur in cesarean birth. Dr. Falls has attempted to overcome this by having an assistant hold the newborn baby by the feet while he gently compresses the chest toward the nasopharnyx, and strips the passages of any mucus. The nasopharnyx is then aspirated.

I believe if a baby is so treated, placed in a crib on one side with the head in a dependent position, and the upper air passages aspirated as frequently as fluid accumulates, the obstructive complications to the newborn delivered by cesarean section will be largely eliminated.

WILLIAM DURWOOD SUGGS, M.D. Richmond, Va.

#### Ménière's Syndrome\*

TO THE EDITORS: Regarding the rationale of vitamin B therapy for Ménière's syndrome, a fundamental medical issue is brought up which every individual physician meets daily. The issue is whether to treat the patient's symptoms or his disease. Ménière's syndrome is not a diag-

nosis but a group of signs and symptoms, and treating symptoms without \*Modern Medicine, Sept. 1, 1949, p. 74.

(Continued on page 98)



#### What do you demand in a toxoid, Doctor?

small dosage volume?
purity?

concentrated potency? high antigenicity?

Of course, doctor, you want the best possible combination of these advantages.

Research and manufacturing know-how of CUTTER, first producer of combined toxoids, have developed in their new purified toxoids products which meet all of your demands. As an example, consider the superiority of the new, purified DIP.PERT-TET\*—for simultaneous immunization against diphtheria, pertussis, and tetanus:

1. Immunization routine is simplified with three injections of 0.5cc each at monthly intervals.

2. Purified toxoids assure virtual freedom from reactions due to bacterial protein components.

3. Allydrox S - CUTTER'S exclusive adsorbing agent - results in a more solid immunity, fewer post-injection reactions, and less pain on injection.

When single immunizations are indicated or booster shots are required, there is a PURIFIED TOXOID—CUTTER—available in both single and multiple dose packages. Your pharmacist has them in stock.

\*DIP-PERT-TET—Cutter's diphtheria and tetanus toxoids and pertussis vaccine combined for simultaneous immunization against diphtheria, pertussis, and tetanus.

\*\*Alhydrox-Trade name for aluminum alhydrox adsorption, exclusive with CUTTER.

**CUTTER** 


CUTTER LABORATORIES . BERKELEY, CALIFORNIA

# in hypertension veratrum viride in CRAW UNITS\*

Of the many drugs commonly used to lower arterial pressure in essential hypertension, veratrum viride (Biologically Standardized in CRAW UNITS) is the only drug that produces a physiologic fall in blood pressure.

A prominent feature in the integrated response to oral doses of veratrum viride in CRAW UNITS is a reduction in peripheral resistance without compromise of circulation and without disrupting circulatory equilibrium.

\* The CRAW UNIT is a multiple of the amount of whole powdered veratrum viride which causes cardiac arrest in the test animal, Daphnia Magna. VERATRITE® and VERTAVIS,® both products of Irwin-Neisler research, are the only therapeutic agents available today that contain a Biologically Standardized veratrum viride in Craw Units.



# Veratrite

Each tabule contains:

Veratrum Viride (Biologically Standardized) 3 CRAW UNITS Phenobarbital.

VERATRITE is a practical preparation for everyday treatment of mild and moderate (Grade I and Grade II) essential hypertension. A calm, gradual fall in blood pressure is assured in the majority of cases, with prompt relief of symptoms. Particularly significant is the prolonged hypotensive response produced by veratrum viride in Craw Units: the fall in blood pressure begins in 1 to 2 hours, reaches a maximum in 4 to 6 hours and ends in 10 to 14 hours.

VERATRITE exhibits a wide range of therapeutic safety, does not require elaborate dosage controls. Supplied in bottles of 100, 500, 1000.

IRWIN, NEISLER & COMPANY (ELINOIS



B LOOD PRESS

Literature and samples on request.

knowing the condition from which they arise is a sure way to court disaster. Unfortunately patients may pay with their lives because of this lazy way of practicing medicine.

It would be well for physicians to attend more clinical pathologic conferences and see unveiled the many conscientious mistakes of the best clinicians who have made earnest efforts to arrive at a diagnosis and even then failed to stem the tide which carried the patient to his death. It should not take any great stretch of the imagination to see how many times these instances of death will be multiplied if we start pumping large amounts of vitamin B into every patient who has Ménière's syndrome and allow the diagnosis or origin of his trouble to go wanting. We know that all excess intake of vitamin B must be excreted, usually unused.

One should never criticize unless he does it constructively and supports his own argument by deeds rather than words. Therefore I refer to my recent article, "The Association of Hyperglycemia, Elevated Blood Pressure, Increased Pulse and Latent Vertical Phoria" (Eye, Ear, Nose & Throat Monthly 28:359-370, 1949), which touches on Ménière's syndrome. Among the patients mentioned in this article are many who have had vitamin B therapy without relief, so at least we will need something other than vitamin B to take care of all who have Ménière's syndrome.

Ménière's syndrome represents a heterogeneous group of cases. The medical profession must ferret out separate causes for each type of case constituting the group. Vitamin B may have a proper place in the partial relief of many, or complete relief of a smaller number, but with the etiology varying from one type to the other, it is expecting too much of any single drug or vitamin to clear up all cases in the syndrome.

I am reminded of the hundreds of patients with pellagra I saw as a boy in Georgia who had vitamin B deficiency in its worst form but who did not have Ménière's. In practice I now see scores of vitamin-B deficiency cases to one of Ménière's syndrome.

I do not wish to detract from any man's effort to bring light where there is darkness, and I am sure this was Dr. Miles Atkinson's purpose in publishing his admirable work with vitamin B therapy. But I dare say that many of his cases will ultimately be found to have an abnormality in blood and body chemistry as the reason for the Ménière-like symptoms. Hyperglycemia is but one of the causative agents, and certainly is not cured by vitamin B therapy.

The work I have submitted does not solve Ménière's syndrome, but I dare say it gets nearer to the basic cause than treating mere symptoms.

E. H. COACHMAN, M.D.

Muskogee, Okla.

#### Correction of Hypospadias\*

TO THE EDITORS: In connection with the article by Drs. Fred Z. Havens and Albert S. Black, in my opinion chordee should be corrected at or about the age of five. Reconstruction of the urethra should be done at or just after puberty.

PAYSON S. ADAMS, M.D.

Omaha

\*Modern Medicine, Sept. 15, 1949, p. 71.

 Now, there is an effective ally against the disease known for its rasping difficulties.

This is Nisulfazole — no recruit, but under trial for eight years — a sulfonamide which differs by carrying a substituted nitro radical. It is given as a suspension, intrarectally, where in relatively high concentration it is in contact with the pathological area.

Of 47 chronic ulcerative colitis patients in an early series treated with Nisulfazole Suspension, 37 could be followed for five years; 34 were then symptom free; three were markedly improved. Some received the

An Advance in the Therapy of

drug for 26 months with frequent blood counts and urinalyses.



Full facts about Nisulfazole sent to physicians on request.

 Major, Ralph H., Am. J. Med. 1:485 (Nov.) 1946.

Nisulfazole®

**Brand of PARANITROSULFATHIAZOLE** 

BREON

Supplied in bottles of: 296 cc. (10 fl. oz.) and 3.78 liters (1 gal.)

George A. Breon & Company

KANSAS CITY MISSOURI RENSSELAER N Y ATLANTA SAN FRANCISCO

#### Retropubic Prostatectomy\*

TO THE EDITORS: The retropubic approach to the prostate gland is a distinct advance in prostatic surgery. Our first experience with this method was in 1945 after reading the technic as suggested by Dr. Terence Millin and hearing him at the meeting of the American Medical Association in Atlantic City and later at the meeting of the American Urological Association in Boston. Since that time. we have done well over 200 operations without a fatality.

The large or medium sized gland is preferred for this approach, the fibrotic type being reserved for the transurethral procedure. We follow the Millin technic altogether, having found it to be most agreeable. The number of days of catheter drainage. and the stay in the hospital have been considerably reduced. The catheter is removed in four to six days and the patient dismissed in ten days or so, healed and able to urinate freely.

The greatest satisfaction has been derived in total retropubic prostatectomy for early carcinoma of the prostate gland, using the Czerny incision. This approach makes an otherwise difficult operation comparatively easy and is less likely to entail surgical error than the perineal approach. The perineal approach has one advantage, however. A section for biopsy can be more readily removed through the perineum, since the posterior lobe is the most common site of carcinoma.

After excising the urethra at the apex of the prostate gland outside the capsule a splendid exposure of \*MODERN MEDICINE, Aug. 15, 1949, p. 67.

the seminal vesicles, vas, and even the ureters can be obtained by elevating the prostate gland at the apex and exposing the posterior portion of the bladder. The seminal vesicles with sheath can be dissected and removed under adequate vision. We now include the sheath. We have not found it necessary to remove a great portion of the bladder or transplant the ureters into the colon, but these procedures can be easily accomplished by the retropubic approach.

The careful identification of the organism and the selection of the proper antibiotic before and after operation provide a smooth and

prompt recovery.

GRAYSON CARROLL, M.D.

St. Louis

TO THE EDITORS: It is difficult to imagine anyone who is proficient at transurethral resection resorting to retropubic exposure, with its wellknown risk of osteitis pubis, for the removal of a sclerotic prostate containing calculi, when the transurethral approach is so much safer and so effective.

The retropubic approach for early cancer is inferior to the perineal in that, with the former approach, one must cut the prostate across and elevate it before the presumably cancerous area can be palpated. From the perineum one comes upon the suspected area in the posterior lobe first-a decided advantage.

Whether the very large benign hypertrophies, or the smaller ones in the presence of stiff hips, strictures. and so forth, should be enucleated supra- or retropubically appears to be



for the Treatment of

# ARTHRITIS and RHEUMATISM

RAYSAL WITH SUCCINATE... The ethical salicylate-succinate formula... Employs three principal ingredients—salicylate, iodine, and succinate... designed to combine the almost specific antiarthritic and antirheumatic action of the salicylates, the stimulating and nutritionally corrective effects of iodine and the salicylate detoxifying action of succinic acid. An ideal companion medication for other therapeutic measures employed in arthritis and rheumatism. RAYSAL WITH SUCCINATE will enhance the efficiency of RAYFORMOSIL... a safe and effective combination for use in your next case. Sample and literature will be sent upon request.

# The Deloxified Salicylate Medicament ENTERIC COATED TABLETS (SALOL)

Available for office use and at your pharmacy on prescription

# RAYMER

PHARMACAL COMPANY • PHILADELPHIA 34, PA. PHARMACEUTICAL MANUFACTURERS

Over a Quarter Century Serving Physicians

a matter of personal taste. There is much to be said for the retropubic approach in these situations, although we need to know more about the incidence of osteitis pubis in ordinary hands.

C. D. CREEVY, M.D.

Minneapolis

▶ TO THE EDITORS: I consider retropubic prostatectomy a definite advance in surgical approach to vesical neck obstructions. Both the mortality and morbidity have been reduced by this method. This approach is in keeping with normal physiologic function of the prostatic method since there is less trauma to the mucosa. Its continuity is better preserved and dissection from adjacent structures to the gland is cleaner and under direct vision.

Hemostasis at times is somewhat difficult, but this was also the "bugbear" of the older methods.

NORVELL BELT, M.D.

Frederick, Md.

# Heparin Therapy of Coronary Thrombosis\*

TO THE EDITORS: Concerning the recent article by Drs. Leo Loewe and H. B. Eiber, on our service we are now using anticoagulants routinely in cases of coronary thrombosis, since in our opinion a cooperative study carried on under the auspices of the American Heart Association proved conclusively that their use has resulted in definite lowering of mortality. Such a lowering has not been apparamount of the state of the transfer of the state of th

ent in a series of 50 patients whom we treated within the past year, but we believe that our series is too small for the statistics to be valid.

We use both heparin and dicumarol, discontinuing the former as soon as the dicumarol has resulted in sufficient reduction of prothrombin activity. Incidentally, we have, during the past two years, been carrying on a controlled study on the use of dicumarol in patients with congestive heart failure, a series now running well over 300 cases. A significant reduction of mortality has occurred, which can be attributed to the practically complete absence of thromboembolic complications in the treated group.

We believe that in a hospital with good laboratory facilities, the only advantage of heparin over dicumarol is its more rapid effect. We believe that if the technics for determining prothrombin activity are carefully standardized, dicumarol can actually be given with greater safety than heparin. If, however, good laboratory facilities are not available, as in the case of general practice in small communities, heparin controlled by determinations of coagulation times is safer, though probably less effective.

EDGAR HULL, M.D.

New Orleans

▶ TO THE EDITORS: In my opinion, anticoagulant therapy is valuable in coronary thrombosis, and dicumarol has been used when no known contraindications existed. I am not able to discuss with authority the use of heparin in coronary thrombosis since I have used it only on occasions to initiate therapy during the period in

# BACITRACIN-NASAL



# More Than Symptomatic Relief IN ACUTE AND CHRONIC SINUSITIS

Bacitracin-Nasal-C.S.C. is a valuable means of reducing the period of disability when acute sinusitis complicates coryza. Bacitracin, through its specific antibiotic properties, destroys many of the pathogens which flourish in the nose and accessory nasal sinuses. Desoxyephedrine, through its vasoconstrictor influence, improves ventilation and sinus drainage, thus enhancing the action of bacitracin. Bacitracin-Nasal-C.S.C. may be administered by means of a nebulizing spray or by the Parkinson lateral head-low position. Available in ½ ounce bottles on prescription at all pharmacies.

- 1. Nonallergenic, even on repeated administration.
- An aqueous solution which does not inhibit ciliary activity.
- 3. Nonirritant, isotonic.

When dispensed by the

pharmacist each cc. of Baci-

tracin-Nasal-C.S.C. provides: bacitracin 250 units,

desoxyephedrine hydro-

chloride 2.5 mg. (0.25%),

sodium benzoate 1%. The solution is stable at room

temperature for 5 to 7 days:

at refrigerator tempera-

ture for 3 to 4 weeks.

4. May be administered to both adults and infants.

C.S.C. Pharmaceuticals

A DIVISION OF COMMERCIAL SOLVENTS CORPORATION, 17 EAST 42ND STREET, NEW YORK 17, N. Y.

which the prothrombin level of dicumarol was being established.

It seems to me that depot heparin offers a chance for anticoagulant therapy to physicians whose work is done in community hospitals where reliable prothrombin level determinations are not available. The cost of prolonged treatment with heparin, however, is still relatively great.

EUGENE H. DRAKE, M.D.

Portland, Me.

#### Liver Function of Chronic Alcoholics\*

TO THE EDITORS: I do not see how one can expect to distinguish liver damage caused by ingestion of alcohol from that due to poor diet or other adverse factors by using laboratory or any other means except the clinical history. After all, so far as we can determine pathologically when we have liver tissue under the microscope, the damage seen in alcoholics is very comparable in all of its stages with that seen in liver damage from a variety of other etiologic causes. Therefore, it is unlikely that any laboratory tests of liver function would show any distinction in alcoholics. In my own experience this has been borne out.

The reports of Drs. Walter L. Voegtlin and William R. Broz comprise a very interesting study on this subject. They were apparently dealing with the early phases of liver damage. In other studies the use of the aspiration needle biopsy has contributed immensely to our knowledge of the pathology in these cases.

DOLPH L. CURB, M.D.

Houston

\*MODERN MEDICINE, Sept. 1, 1949, p. 50.

► TO THE EDITORS: When one starts reviewing opinions of others on a controversial subject, confusion results. So it is in this case. Even as far back as the early 1930's, when it was almost axiomatic that a cirrhotic liver was due to alcoholism, doubters of this theory began appearing.

It is my opinion that a great majority of investigators now believe that liver damage is not due to alcohol per se but to dietary insufficiency. Fatty infiltration of the liver is the precursor of cirrhosis, and this may be caused by dietary insufficiency, deficiency diseases, chronic ulcerative colitis, diabetes mellitus, or pigment as seen in hemachromatosis. Withdrawal of dextrose from body metabolism may result in fatty liver.

Production of cirrhosis with alcohol experimentally has not been successful. However, as one authority states, if cirrhosis is suspected, inquiry usually is made concerning the use of alcohol; if alcoholism is suspected, a diligent search for cirrhosis will be made. Aside from this, little, if any attention is given the matter. Case history studies are most unreliable, except when the two conditions are under investigation.

It would be unfortunate, however, for the layman to gain the impression that alcohol is blameless in the development of cirrhosis; certainly it must be much to blame and given substance. All in all, the decision is not closed, but I feel that the consensus now is that liver damage from ingestion of alcohol cannot be distinguished from that due to poor diet and other adverse factors.

MORRILL L. ILSLEY, M.D. Claremont, Calif.

individuality
in
spasmolytic
prescription
writing

Individualized management of complex spastic dromes is particularly facilitated by use of Donno Elixir—the spoonfed spasmolytic agent whose superior efficacy derives from its precise balance of the principal natural alkaloids of belladanna, plus phenobarbital • With it may be administered judicious selections from a wide variety of gastric, intestinal, branchial, urinary, analgesic, or other drug agents with which it is entirely compatible for the precise pharmacotherapy you desire.

Each Sec of Donnatal Blinic contains:

Hyoscyamine Sulfate		
Atropine Sulfate	,	
Hyossina Hydrobromi		
Phonoburbital (% gr.)	ILS mg	

spusmolysis plus

donnatal elixir

Also available as

Donnatal Tablets

and Donnatal Capsules



A. H. Robins Co., inc.

Third Pharmaceuticals of Marit since 1978
Richmond 20, Va.



the synergic formula for maximum non-narcotic analgesia

To obtund pain without recourse to narcotics-yet better than the patient's medicine cabinet can-becomes a daily professional obligation. That's why Phenaphen was formulated with calculated pharmacologic precision . . . the analgesic action of its aspirin - phenacetin components being implemented and prolonged by its phenobarbital content (which helps allay apprehension . . . its hyoscyamine further increasing overall efficiency through local anodyne action. Phenaphen -the astute professional prescription for pain - is promoted to physicians only.

#### Each tablet or capsule contains:

Phenacetin (3 gr.)	194	mg.
Acetylsalicylic Acid (21/2 gr.)	162	mg.
Hyoscyamine Sulfate	0.03	mg.
Phenoharbital (1/4 er.)	16.2	****

• phenaphen Robins



A. H. Robins Co., inc. ETHICAL PHARMACEUTICALS OF MERIT SINCE 1878 Richmond 20, Va.

▶ TO THE EDITORS: Ethyl alcohol has never been firmly established as the direct etiology of the so-called "alcoholic cirrhosis," but there is no question that by inference it has been implicated as a contributory factor. Approximately 50% of all patients with Laennec's cirrhosis are heavy drinkers, but by the same token, in countries with a large population of Mohammedans, who are abstainers, the incidence of cirrhosis is equal to that of the United States.

The present concept points to alcoholism as a cause of poor dietary habits and alterations in metabolism of essential food elements and vitamins, this nutritional deficiency in turn being the direct cause of the

liver pathology.

There is little argument that in the late stages of cirrhosis, namely the small hobnail liver, nothing in the pathologic picture will indicate whether the patient was or was not alcoholic. However, there is some evidence that in the earliest stages of liver involvement specific cytologic changes are present which may be attributable to ethyl alcohol.

Mallory, in 1933, reported the presence of an acidophilic hyaline reticulum in the hepatic cells of the central portion of the lobule, characteristic of the earliest phase of alcoholic cirrhosis. This initial lesion precedes the development of portal fibrosis. However, he was later able to produce the identical lesion by experimental phosphorus poisoning.

Hall and Morgan have pointed out that there is a very specific subacute alcoholic cirrhosis which may be differentiated pathologically from other forms of liver disease. Its differential points include a more cellular type of portal connective tissue proliferation, a more widespread necrosis of hepatic cells, and again, as observed by Mallory, a hyaline necrosis of the cytoplasm.

It must be remembered that the cases of Hall and Morgan were in the subacute stage of the disease and that this picture is comparable to that of early toxic cirrhosis due to other causes than alcohol. From a clinical standpoint these observations are purely of academic interest, since pathologic observations in the very early stages of the disease are not usually within the realm of the physician, and laboratory investigation cannot distinguish the exact type of cellular lesion.

At the present time, for all practical purposes, there is no method by which liver disease associated with chronic alcoholism can be distinguished from hepatitis and cirrhosis due to other causes.

SIDNEY A. PORTIS, M.D.

Chicago

## Giant Cell Tumor of Bone\*

TO THE EDITORS: Dr. T. M. Prossor's article refers particularly to osteoclastoma. It is accepted that this tumor is best treated by radiotherapy. Only bone tumors that are known to be radiosensitive can be so treated. Unfortunately, most bone tumors are not radiosensitive and require surgery.

ABRAHAM MYERS, M.D.

Philadelphia

\*MODERN MEDICINE, Oct. 1, 1949, p. 81.

► TO THE EDITORS: The relative merits of radiologic and surgical treatment for bone tumors cover a rather wide and unsettled field, but certain principles can be enunciated from the conjoint experience of the many who have made comprehensive studies in it. There are certain instances in which radiologic treatment holds sway, and certain ones in which surgery is the treatment of choice.

When any bone tumor is quite accessible, there can be no question—complete excision is the method of choice. This applies to both benign

and malignant tumors.

In osteogenic sarcoma, the exact surgical procedure depends upon the degree of malignancy. If it is grade 1, more conservative surgery is indicated—in grade 4, amputation is the method of choice. This is evidenced by the number of five-year cures in published series of cases.

In fibrosarcoma, early amputation

offers the best prognosis.

Giant cell tumor of bone is best treated by surgery; the exact type of operation depends upon the location of the lesion. Complete excision when possible is always the method of choice. Inaccessible lesions are treated by radiologic method only as second choice. In malignant giant cell sarcoma, amputation gives the best prognosis.

Of the vascular tumors of bone, accessible benign hemangiomas are treated surgically if treatment is required. If not accessible, radiologic treatment is indicated. Since benign hemangiomas rarely give much trouble, too great risk is not being taken. On the other hand, in Ewing's tumor or malignant angioma or en-

dothelioma the treatment of choice is radiologic.

Multiple myeloma is not affected by any form of treatment, surgical or radiologic.

M. E. PUSITZ, M.D.

Topeka, Kan.

## Acute Rheumatic Disease in Children\*

TO THE EDITORS: Since we know that at least 50% of adult patients with definite rheumatic endocarditis give no recognizable history to suggest rheumatic infection, the comments of Dr. Leo M. Taran are timely. He points out that acute rheumatic heart disease is not a clear-cut entity, but it seems most unwise to cast aside laboratory evidence.

While acute carditis may occur without elevation of the sedimentation rate, white blood cell count, temperature, pulse rate, or antistreptolysin, such occurrence is so rare that perhaps it is unwise to stress the

point.

In average practice one sees so much unwarranted cardiac invalidism that it is felt that the practitioner must avail himself of as much concrete evidence as possible with definite associated cardiac findings. Only then will the incidence of functional murmurs treated as organic be reduced. Occasional organic murmurs are going to be missed for a few years, but this is certainly a lesser evil than the tragedy of a child saddled for life with the diagnosis of heart disease because of a functional murmur.

G. I. BELL, M.D.

Edmonton, Alb.

\*Modern Medicine, Feb. 15, 1949, p. 55.

OF CROSS
INFECTION
BACTERIA
WITH





Recent controlled investigations have confirmed the low concentration effectiveness of triethylene glycol vapor against airborne streptococci, pneumococci, staphylococci and other disease-producing organisms.

**NOW**—this principle has been made practical, simple and economical for use in your office, the hospital, the home. One spraying—only a few seconds—is sufficient to reduce airborne bacteria as much as 90%.

- Vapor is quickly dispersed to all parts of room.
- Nontoxic, odorless, harmless to furnishings and equipment.
- Avoids cumbersome, expensive apparatus.
- Costs only a few cents a day.
   Write for information on "MICROBOMB"

# THIS IS THE TRADEMARK OF THE CARAND CORPORATION

CARAND CORPORATION . RACINE, WISCONSIN

## Diagnostix

Here are diagnostic challenges presented as they confront the consultant from the first clue to the pathologic report. Diagnosis from the Clue requires unusual acumen and luck; from Part II, perspicacity; from Part III, discernment.

#### Case MM-155

#### THE CLUE

ATTENDING M.D: The young woman in the next room is twenty-nine years of age and has had abdominal complaints for a long time and many operations. In the past nine years she has had uterine suspension, salpingectomy, drainage of a right polycystic ovary, cholecystectomy, two dilations and curettages, and amputation of the uterine cervix. Now the surgeons want to do a laparotomy.



visiting M.B: The operations have apparently been of little value. This is about the time that one begins to suspect a psychiatric or at least an obscure medical condition. Let us assume that it's the latter to start with. What clues can you give me?

has had a palpable mass in the left lower quadrant. It is tender and quite large—perhaps 15 by 10 cm. During an exploratory operation eight years ago, the left ovary was found to contain numerous cysts, and it was drained and marsupialized. We have no details of this operation. For six years the patient has had recurring attacks of bilateral costovertebral pain radiating to the epigastrium and, during these spells, abdominal tenderness and rigidity.

visiting M.D. I presume nothing was found at these operations to account for her symptoms; at least, she apparently still has attacks of pain.

ATTENDING M.D. Correct.

#### PART II

ATTENDING M.D: She has been hospitalized now because of the abdominal mass, tenderness, rigidity, nausea, vomiting, and an intermittent, cramp-like pain, which is intensified just before a bowel movement.

# The Advantages of Vicks VapoRub in Steam

#### WHEN STEAM INHALATION IS INDICATED

Whenever you recommend steam therapy for the relief of respiratory distress, your patients will find Vicks VapoRub in steam especially effective in bringing prompt, soothing relief.

VapoRub's well-balanced formula contains 7 volatilizing ingredients including menthol, thymol, camphor, and oil of eucalyptus.

Used in steam, VapoRub provides soothing vapors that are comforting to a patient—particularly when dryness and irritation of the mucous membranes are



IN VAPORIZER



IN BOWL

prominent features accompanying the respiratory infection.

VapoRub can be used in a vaporizer or a bowl of boiling water. It can be used as often as needed—for all the family.

Vicks VapoRub is available in practically every home. In recommending it, you can be sure that it's already on hand for instant use. When steam inhalation is prescribed, recommend . . . Vicks VapoRub.

She also has diarrhea with twenty to thirty soft stools a day.

VISITING M.D: (Walking into the room. The patient has a Wangensteen tube in the bowel and is getting a blood transfusion) Now, what immediately preceded the symptoms you have described?

ATTENDING M.D: For six months the patient has had scanty, irregular menses; for three weeks the abdominal mass has been growing more sensitive and enlarging. For two weeks she has had unbearable abdominal pain. Today there is some rebound tenderness, greater on the left.

VISITING M.D.: Do the laboratory tests help us?

ATTENDING M.D: She has had the works: from blood urea and amylase to urine porphyrin. Results of all the special tests are negative and those of the routine studies can be accounted for on the basis of dehydration or mild infection. The sedimentation rate is 70. She has lost only 10 lb. in the past five years. VISITING M.D: How about films of chest

#### PART III

and abdomen?

ATTENDING M.D: The chest film is negative. The roentgenogram of the abdomen reveals numerous cystic calcified areas measuring about 0.2 to 1.3 cm. They are round, smooth, discrete, are more dense peripherally, and some appear lobulated. All other x-ray studies, including excretory urograms, are negative.

VISITING M.D. Mmmm . . . Young woman, recurrent episodes, ovarian cysts, abdominal calcification. (To himself) Paraffinoma, echinococcus

cysts. calcified mesenteric nodes. phleboliths, biliary or urinary calculi, calcified fibroids, enteroliths, calcified primary or metastatic tumor . . . pseudomyxoma peritonaei . . partial bowel obstruction. (Aloud) I'd like to examine the patient proctoscopically. (After examination) I note some small rectal nodules and a stricture-like shelf on the anterior rectal wall. I believe this is pseudomyxoma peritonaei. I'd advise peritoneoscopy before surgery, but would continue conservative therapy for bowel obstruction until things are quieter. Of course, it may be necessary to explore, but I'd hesitate to do that now.

#### PART IV

ATTENDING M.D: (One week later) The patient responded well and peritoneoscopy and biopsy confirmed

your opinion.

VISITING M.D: Undoubtedly she will continue to have episodes of partial bowel obstruction. These nodules and tumor-like growths usually result from rupture of a mucocele of the appendix or pseudomucinous cystadenoma of the ovary. The pseudomucinous material and the cells producing it are spread through the abdominal cavity. The peritoneum reacts with chronic inflammatory changes, and adhesions are formed. There may be proliferation of the implanted cells with small, tumor-like growths. Some believe the mucoid material produces the lesions, but I doubt it, although the material is undoubtedly an irritant and provokes a foreign-body reaction.

## It is a Question of Management..

The treatment of obesity is simple. It is the management of the obese patient that presents the problem.



DIETENE® is of real value in the management of the obese patient, particularly during the early days of diet restrictions, when the pangs of hunger may tempt the patient to deviate from the regimen.

DIETENE supplies essential protein, vitamins, and minerals in a very palatable form. One-pound cans, plain or chocolate flavor, are available to patients through all pharmacies at \$1.55.

Send for your free supply of the 1000-calorie Dietene Reducing Diets. Diets are made up to look as if they were typed in your office for the individual patient.

Specify DIETENE in reducing diets to maintain adequate protein-vitamin-mineral intake.



Not advertised to the laity

THE DIETENE COMPANY 518 FIFTH AVENUE SOUTH, MINNEAPOLE	Dept. DM 119 5 15, MINN.
Please send me a free supply of the Diet.	1000-calorie Dietene Reducing
Name	
Address	
CityZor	eState

### Interpreting Medicine for the Layman

STEVEN M. SPENCER\*

Associate Editor, The Saturday Evening Post

The private practice of medicine is much more medicine and much less private than it was ten or fifteen years ago. The doctor, striving to familiarize himself with and to evaluate the parade of new drugs and technics, finds an eager public looking over his shoulder—and sometimes breathing down the back of his neck.

Gone are the days when a practitioner could keep the details of diagnosis to himself and dismiss Mrs. Brown with a reassuring word to the effect that "it's nothing serious—and if it is come back on Friday and we'll try something else." Mrs. Brown will want to know the name of the germ that bit her. She will also ask if the prescription contains a sulfa, which upset her digestion last time she took it, or penicillin, which upset her budget.

However much you may be annoyed by the sound of a little medical knowledge rattling around in a patient's skull, you probably will agree that the public has a legitimate interest in your problems and progress. Steven M. Spencer believes that this interest can be a healthy situation not only for the patient but for the doctor and his profession as well. People who fully understand and appreciate what the great American medical profession has done and is continuing to

do are less likely to accept a compulsory health insurance scheme or any other socialized medicine package wrapped in Washington and tied with yards of red tape.

Much of the public's information about medicine is obtained through the press. If that information is sound the public reactions on medical matters are more apt to be sound than otherwise.

Medicine is not something people take instinctively, as they take food or water. It is a commodity on which they must first be sold. Selling is largely a matter of telling. That is where physicians and writers and editors can work together. For it seems logical that the more people know about what doctors can do to help them, the more readily will they come to doctors for help, and the more readily will they rally to the doctor's point of view in controversies over medical economics.

Although most people take medicine only when they or their physicians feel it is needed, information about medicine can be ingested, without discomfort, at almost any time. For the human machine has ever been a fascinating subject.

A popular interest in medicine has always existed, even though the doctors and editors may have underestimated it in the past. One of the first

## eradicate scabies

WITHIN 24 HOURS

# EURAX

BRAND OF CROTAMITON CREAM (contains 10% N-ethyl-o-crotonotoluide)

# scabicidal non-irritating antipruritic bacteriostatic

Clinical experience has demonstrated that a single application of EURAX cream completely eradicates scabies in more than 88 per cent of cases. Two applications, 24 hours apart, produce cure-rates up to 100 per cent.  $^{1.6}$ 

EURAX - "... free from many, possibly all, of the objectionable features of other sarcopticides" - a totally new product of Geigy research, offers the following unique advantages:

Prompt relief of pruritus is almost universally obtained within an hour of application.

Associated pyoderma is remarkably alleviated. No preliminary scrubbing necessary.

Non-irritating, non-toxic in recommended dosage. Non-greasy, non-staining, and non-odorous.

EURAX (Brand of crotamiton): Available in 10% concentration in a vanishing cream base, tubes of 60 Grams.

BIBLIOGRAPHY: 1. Couperus, M.: J. Invest, Dermat. 13:35, 1949. 2. Tronstein, A. J.: Ohio State M. J., in press, 3. Goldman, L.: Connecticut M. J. 13:62, 1949. 4. Patterson, R. L.: Work in progress, 5. Domenjos, R.: Schweiz, med. Wchnschr, 76:1210, 1946. 6. Burckhardt, W., and Rymarowicz, R.: Schweiz, med. Wchnschr, 76:1213, 1940.





The world wide Geigy organization—established in Switzerland in 1764—has a noteworthy history in the production of synthetic organic compounds which have found wide acceptance as chemicals of highest quality. Leadership in organic research and synthesis have led naturally to the development of original pharmaceuticals now widely prescribed in virtually every country of the world.

GEIGY COMPANY, INC. 89-91 Barclay St., N. Y. 8, N. Y.

newspapers printed in Europe carried in 1494 an account of the outbreak of syphilis in Naples following the return of Columbus and his men from the New World. Every new development that comes over the medical horizon is greeted with a fresh spurt of printer's ink. The sulfa drugs were one of the big stories of the century. Penicillin, the discovery and application of which were filled with drama, made an even greater impact on the people's imagination.

Most medical news is good news. It has been said that the good and the peaceful make dull reading. But in this day when so much of the news is bad, good news is doubly welcome. And medicine is not dull at any time. Even when an article does not an-

nounce a brand new treatment but simply presents a summary of the most up-to-date knowledge on the cause and treatment of any disease, or reports a promising development along the research front, it gets a good reading. For the reassurance which comes from a fuller understanding of a condition affecting the reader or a member of his family is itself ample justification for publishing the information.

A case in point was the article, "I Have a Scar on My Heart," a personal experience story by a Detroit advertising man, W. A. P. John, which the Saturday Evening Post printed last winter. It did not announce a sensational new therapy. It was just a well-written account of how the

## Accepted by Leading Pediatricians Accepted by Council on Foods

Laxative Modifier of
Milk for Constipation
in Infants

Borcherdt
MALT SOUP EXTRACT

DRI-MALT SOUP EXTRACT
(Powder)

(Liquid)

FEATURES:

 High proportion of readily fermentable maltose encourages the growth of aciduric bacteria and retards growth of putrefactive organisms.

Malt Soup Extract stimulates peristalsis.
 Water-soluble extractives of choice, malted barley and the added potassium carbonate contribute to the gentle laxative effect.

Mixture of sugars (maltose-dextrins) means better toleration—no danger of gastrointestinal irritation, excessive fermentation, or diarrhea when used as directed by the physician.

Palatable... Dissolves Readily in Milk SUPPLIED: Malt Soup Extract—Jars containing 8 fl. oz. and 1 pt. Dri-Malt Soup Extract—Jars containing 1 lb.

Borcherdt MALT EXTRACT COMPANY
Malt Products for the Medical Profession Since 1868
217 NORTH WOLCOTT AVENUE CHICAGO 12, ILLINOIS

#### CHECK LIST

## for choice of a laxative

## Phospho-TYPE OF Soda (FLEET)\* ACTION

- ✔ Prompt action
- ✓ Thorough action
- ✓ Gentle action

#### SIDE EFFECTS

- Free from Mucosal Irritation
- Absence of Constipation Rebound
- No Development of Tolerance
- Safe from Excessive Dehydration
- No Disturbance of Absorption of Nutritive Elements
- Causes no Pelvic Congestion
- No Patient Discomfort
- ✓ Nonhabituating
- Free from Cumulative Effects

#### ADMINIS-TRATION

- Flexible Dosage
- Uniform Potency
- ✔ Pleasant Taste

### Judicious Laxation



## through controlled action

Phospho-Soda (Fleet)\*, over the years, has won discriminating preference by thousands of physicians . . . because of its controlled action — its freedom from undesirable side effect—and its ease of administration. Your prescription of Phospho-Soda (Fleet)\* assures effective (and safe) results. Liberal samples on request.

#### C. B. FLEET CO. INC.

ITNCHBURG VIRGINIA

PHOISPHO SODA and FIEES

reduction as or C B. Fleer Cb., Inc

PHOSPHO-SODA

Phospho-Seda (Fleet) is a solution containing in each 100 cr. (addum

CCEPTED FOR ADVERTISING BY THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION

writer had come through a coronary attack, how he had taken a philosophic view of the situation, how he had slowed down his pace, how, in brief, he was following his doctor's orders and getting along fine.

The response to this article was overwhelming. Mr. John received nearly 500 letters within a month after publication, virtually all of them commendatory and many expressing gratitude for some individual help or encouragment the reader had derived. Several wives thanked Mr. John for having dramatized the dangers of overdoing, and said their husbands had at last begun to heed the warning, even though up to that time they had completely disregarded the same advice from their own doctors.

This is one kind of medical information that is helpful to the patient. If it makes the doctor's job any easier it may be considered helpful to him, too.

Education of the public in the value of early diagnosis of cancer has now reached the point where the demand for examinations exceeds the facilities for conducting them. Dr. Elise L'Esperance, Director of the Strang Cancer Prevention Clinic in New York, has stated that there is a waiting list of 6,000 in that area alone, and that a comparable situation exisits in Philadelphia, Pittsburgh, Chicago, and other cities where cancer detection centers are operated.

Now a layman who is interested in

# FOR ECZEMA

"—the advantage of the diminution of the black color is obvious"\*

# SUPERTAH (NASON'S) WHITE, NON-STAINING DINTMENT Has Other Advantages:

An authoritative work on skin diseases says of SUPERTAH: "It has proven as valuable as the black coal tar preparation . . . it does not stain the skin or clothing, nor does it burn or irritate the skin.

\*Swartz & Reilly, "Diagnosis and Treatment of Skin Diseases," p. 66.



It can remain on the skin indefinitely without fear of dermatitis."\*

SUPERTAH (Nason's) is a white creamy ointment, packaged in original 2-oz. jars, 5% & 10% strengths. Distributed ethically.

TAILBY- NASON COMPANY
Kendall Square Station, BOSTON 42, MASS.

# On the Dual Value of

Socially candy has long been accepted as a pleasant part of our daily lives. From early childhood on, candy is considered an appropriate accompaniment of the festive spirit of birthdays, holidays, anniversaries, and

In recent years, with the advancing knowledge of nutrition, the values of candy as a worth-while part of the daily diet have also become recognized. Most of the kinds of candy manufactured today are made of a number of valuable foods which contribute to the extent they are used to the satisfaction of many nutritional needs.\*

other joyous occasions.

Whether enjoyed as a delectable tidbit during a friendly gathering—or served at the end of a family meal—or eaten as a quick energy food following strenuous activity, candy has a unique and valid place in the human dietary.

"The candies in the manufacture of which milk, butter, eggs, fruits, nuts or peanuts are used, to this extent also (a) provide biologically adequate proteins and fats rich in the unsaturated fatty acids; (b) present appreciable amounts of the important minerals calcium, phosphorus, and iron; (c) contribute the niacin, and the small amounts of thiamine and riboflavin, contained in these ingredients.

Council on Candy

OF THE

NATIONAL CONFECTIONERS' ASSOCIATION
ONE NORTH LA SALLE STREET, CHICAGO 2, ILLINOIS

medical details often expects more information from his doctor than the latter may feel he has time to supply. Here again the press can perhaps be of help to both the physicians and the patients.

For many readers magazine articles answer questions which have long bothered them about the general nature of their ailments. Specific details about a patient's own case obviously can be supplied only by his doctor. But through the medium of the press, a type of background information can be provided that in the long run will save you an endless amount of time and your patients an endless amount of worry.

It is more and more to the advantage of the medical profession today

to meet the public halfway on these matters, for it is the public to which the doctors must look for an increasing share of support for medical education and research. As taxpayers providing government grants and as voluntary contributors to a growing list of special campaigns, the people want to be kept informed on medical affairs and they want to be assured that the doctors examine both sides of medical economics questions. If the American medical profession is to have an electorate favorably disposed toward the physicians' views on compulsory health insurance, for example, that electorate must be kept sold on the American medical profession.

The story of American medicine

# for **RELIEF** of constipation without catharsis

# neo-cultol®

Natural Corrective Non-Habit Forming

Arlington

THE ARLINGTON CHEMICAL COMPANY

YONKERS 1, NEW YORK

L. acidophilus in refined mineral oil jelly, chocolate flavored—restores normal intestinal flora and normal colonic function without griping. flatulence, diarrheic movements—gently lubricates without leakage. Jars containing 6 oz.



\*ALSO KNOWN AS DETTOL

# A proved antiseptic for obstetrical and surgical use

• Dett, known as Dettol throughout the British Empire and other parts of the world, is now available to the medical profession of the United States.

Dett, although deadly to germs, is gentle to human tissue. This clean, clear liquid with an agreeable odor is safe, effective, non-

irritating and non-staining. Physicians who have used Dettol in other countries will welcome its introduction in the United States under the name of Dett.

For a generous size sample, and literature, write to: The R. T. French Co., Pharmaceutical Dept., Rochester 9, New York.

DETT THE MODERN WEAPON AGAINST INFECTION

is living, contemporary history. As such it must concern itself with living men and women. Medical contributions are not made by disembodied spirits. They are made by individuals and teams of individuals-research workers in their laboratories, surgeons in their operating rooms, general practitioners and specialists in their offices and at the bedside. And the light of the whole group will shine but dimly, so far as the public eye is concerned, if the light of each member of the group is hidden under his own private bushel of anonymity. The history of medicine would be dull reading indeed if it contained no mention of the names and personalities of men like Harvey, Jenner,

CHIR OPODIST

Beaumont, Pasteur, Osler, Cannon, Cushing, Fleming.

The argument for using names is simply that the public has a right to know whose work is being described and whose opinions are reflected in the text of the article. To adopt a policy of not quoting authorities would open the way for unreliable reports on medicine by irresponsible publications—and there are a few. In addition, we feel the story of medicine's advance is the story of people as well as of facts.

The tradition of medical modesty apparently is based on the theory that if it were declared ethical to blow one's own horn, the quacks would blow loudest, attract most of the business, and ruin the public's health. The theory is all right, but the fact of the matter is that the quacks disregard all rules against hornblowing and do pretty well for themselves. Self-advertising, however, should be clearly differentiated from bona fide medical news. Accurate articles on medical progress are useful and writers should be able to obtain information from doctors without fear of subjecting the latter to censure by their colleagues.

A Code of Cooperation already has been adopted by the Colorado State Medical Society and the press and radio of that state.

The code sets up a system of official spokesmen for each county society—usually the president, secretary, and publicity chairman. These men and women, as well as spokesmen for the hospitals, are to make themselves available to the press and may be quoted "in matters of public interest for purposes of authenticating infor-





The fluid that inundates the tissues during congestive heart failure may pass through approximately one and one-half acres of capillary wall. Following an intramuscular or intravenous injection of MERCUHYDRIN, edema fluid comprised of water and salts, chiefly sodium chloride, is mobilized back through the one and one-half acres of the capillary bed and is eliminated through the kidneys. The diuresis obtained with MERCUHYDRIN benefits not only the patient with palpable edema, but also the patient subject to cardiac decompensation. "The effect on dyspnea in these cases of left-sided failure is probably largely a result of diminution in pulmonary edema, even though the latter is clinically occult."\*

The management of cardiac decompensation is greatly facilitated and the comfort and well being of the patient is greatly increased by administration of

MERCUHYDRIN early, concurrently with digitalization

MERCUHYDRIN in a systematic schedule of repeated doses as maintenance therapy

MERCUHYDRIN by intramuscular injection, well tolerated locally and systemically, and affording highly effective diversis

MERCUHYDRIN (meralluride sodium) is available in 1 cc. and 2 cc. ampuls

\*Fishberg, A. M.: Heart Failure, Lea and Febiger, Phila., 1946, p. 733.



mation." The code states specifically that this action by the spokesmen "shall not be considered by their colleagues as a breach of the time-honored practice of physicians to avoid personal publicity, since it is done in the best interests of the public and the profession." That single sentence cuts to the very core of the problem.

Colorado's program is a tremendously encouraging sign that the physicians and the press are alive to the importance of giving accurate medical information to the public. If the code works out, plans of this type, perhaps broadened in some respects. may be adopted in other states.

For the story of medicine is one which the press and the public are


eager to hear. Your town wants to know what kind of medicine it has, what facilities and services are available, what types of operation its surgeons are performing, what research projects are under way. The public wants to know what you are doing and saying. Your story and every doctor's story is part of the fabric of its life. It is the story of big and little miracles, of birth and growth and struggle and triumph. It is the story of man himself, and that is still the biggest and best story on earth.



For Effective Treatment of ....

## HYPERTENSION

Tablets VERUTAL (Rand)



Prolonged Vasadilation Capillary Protection Mild Sedation Therapeutic Safety

Verutal Tablets (Rand) combine four therapeutically effective drugs in a new formula for the treatment of Essential Hypertension

AND

PHARMACEUTICAL CO., INC.



Tyree's Antiseptic Powder offers the busy physician a balanced vaginal douche . . .

**BALANCED** Psychologically . . . by imparting immediately a sense of cool, clean, gratifying comfort, Tyree's restores the woman patient's subjective balance and makes her amenable to further curative treatment.

BALANCED Physiologically . . . by correcting hypo-acidity present in the vaginal pathology with Tyree's, it is possible to approximate the normal vaginal pH of 4.0—a condition very hostile to the growth of vaginal infections.

BALANCED Therapeutically . . . finally, Tyree's value as a vaginal douche is positive, because it balances effectiveness with safety, avoids complications caused by caustic, irritating douching, while it acts as an effective treatment in vaginal infection. Try Tyree's the next time you prescribe a vaginal douche. Write for literature and professional samples.

## Tyree's ANTISEPTIC POWDER

J. S. TYREE, CHEMIST, INC., 15th and H Streets, N. E., Washington 2, D. C.

Manufacturers of CYSTODYNE, Tyree, for the treatment of genito-urinary infections

HANDY TIMESAVERS

#### FOR YOU!

#### PEDIATRIC FEEDING DIRECTIONS

(birth to 3 mos., 3-6 mos., 6-10 mos., over 10 mos.)

Easy to use, complete, adaptable to individual patients. Help mothers follow your directions accurately. Each contains: for-

mula or diet charts; food lists; food preparation methods; weight record; spaces for your directions, next appointment. Available in pads of 50 each, imprinted if desired.





Eight-page book with pictures for the youngster to color. Written in primer style. Emphasizes health practices and other good habits you and the mother want the child to develop. Yours—to give your young patients.

Use this coupon for sample copies and a postage-paid order card. After examination, mail card to order in quantity.

#### NO COST OR OBLIGATION!

RALSTON PURINA COMPANY, Nutrition Service 9M-2Checkerboard Square, St. Louis 2, Mo.

Please send the FREE material checked below:

C848 1 set Feeding Direction Forms

C958 1 child's Color Book

Name M. D.

Street\_\_\_\_\_

City\_

# When Rapid Growth Calls For HIGH IRON and THIAMINE— Remember RALSTON




instant Raiston — enriched whole wheat cereal—is a *rich* source

of iron and thiamine. A single 1-ounce serving supplies the following percentages of the minimum daily requirement:

1-6 years 6-12 years adults
IRON 113% 84.9% 84.9%
THIAMINE 84% 56% 42%
Plus 3.5 Gm. PROTEIN



Cooks in 10 seconds







## Short Reports

INDUSTRIAL MEDICINE

#### Mercury Poisoning from Fingerprint Photography

A new form of a long-established occupational hazard, chronic mercurialism, is reported by Drs. John N. Agate and Monamy Buckell of London. Policemen who develop latent fingerprints are the victims. In preparation for photographing prints left at the scene of a crime, a powder is dusted over the area. The most commonly used powder by British and American police forces is mercury with chalk. Exposures in excess of two hundred and fifty hours a year are considered to constitute a definite risk. Examination of 32 policemen of the Lancashire constabulary regularly engaged in developing latent prints revealed 7 who had chronic mercury poisoning. The three main symptoms are stomatitis, tremor, and excessive irritability.

Lancet 257:451-454, 1949.

NEUROPSYCHIATRY

#### Alpha Rhythm of the Brain

Voluntary muscular movements are apparently influenced by the electrical rhythms of the brain. Dr. G. O. Kibbler and associates of Whitchurch Hospital, Cardiff, Wales, made simultaneous recordings of the alpha rhythm and eye opening in response to an auditory signal and measured the time interval between the two. The eye tended to open at the peak of the rhythm.

Nature 164:371, 1949.

TREATMENT

#### Relief of Radiation Sickness

Dramamine is effective, prophylactically and therapeutically, against radiation sickness. A total of 300 mg. is given in three equal dosages: thirty to sixty minutes before radiation, an hour to an hour and a half after, and three hours after. Dr. John W. Beeler and associates of the Mayo Clinic, Rochester, Minn., report that the relief of symptoms was significantly greater from Dramamine than from placebos. Drowsiness is the most noticeable side effect. A combination of Dramamine and pyridoxine may be desirable for the treatment of severe cases.

Proc. Staff Meet., Mayo Clin. 24:477-483, 1949.

HORMONES

## Effect of Cortisone on Cerebral Activity

In addition to relieving pain, stiffness, and disability of patients with rheumatoid arthritis, cortisone apparently acts as a mental stimulant. Drs. Edward W. Boland and Nathan E. Headley of the University of Southern California, Los Angeles, made electroencephalographic studies before and after administration of the hormone in 2 cases. In both instances tracings made at the termination of cortisone acetate administration revealed increases in frequency of the alpha waves. In one case the increase was 8.8 and in the other 9.2%.

J.A.M.A. 141:301-308, 1949.

## "The Favorite Medication"\*

## Among Antihistaminics Used in the Pioneering Study AGAINST THE COMMON COLD

\*"Neo-Antergan was found to have little or no sedative effect in the majority of cases in which it was given, and thus became the favorite medication of the ambulatory patients who had had experience with any of the other antihistaminic drugs."

> —Brewster, J. M., Antihistaminic drugs in the therapy of the common cold, U. S. Naval Medical Bulletin 49:1-11, Jan.—Feb. 1949.

Neo-Antergan†—the favorite antihistaminic—is characterized by high antihistaminic potency and a high index of safety. It may be administered with confidence whenever safe, effective antihistaminic action is desired.



†Neo-Antergan is the registered trade-mark of Merck & Co., Inc. for its brand of pyranisamine.



MERCK & CO., Inc. Manufacturing Chemists RAHWAY, N. J.

# Neo-Antergan®

(Brand of Pyranisamine Maleate)

(N-p-methoxybenzyl-N', N'-dimethyl-N-a-pyridylethylenediamine maleate)




DIAGNOSIS

#### Diffuse Pulmonary Lesions Differentiated by Biopsy

When conventional methods fail to establish the nature of diffuse pulmonary lesions, biopsy may permit a rapid, accurate diagnosis. Lung biopsy has been used in 50 cases by Dr. Karl P. Klassen and associates of Ohio State University, Columbus, to diagnose such conditions as tuber-culosis, histoplasmosis, pneumoconio-

causes the lung to herniate through the incision. The edge of the lobe is grasped by a small lung clamp and a mattress suture of 000 chromic surgical gut with a swaged curved needle at each end is placed at the apex of the biopsy site, 2 cm. from the lung margin. With two hemostats approximating the suture, the isolated wedge of pulmonary tissue is excised, c. The two ends of the suture are then run down, as an over-and-over stitch, to



sis, sarcoidosis, and cancer. The procedure is well tolerated and does not induce pneumothorax or hemothorax. Dissemination does not occur in instances of bacterial or fungous diseases. With patient supine, an 8-cm. incision, a, is made over the fourth anterior intercostal space, 4 cm. from the edge of the sternum. Blades of a medium-sized Richardson retractor inserted parallel with the ribs and rotated 90 degrees outward, b, spread intercostal space to expose the interlobar fissure. Slight increase of intrabronchial pressure by the anesthetist

the periphery of the lung on each side, d, the clamps are removed, and suture is tightened and tied, e. One end of the suture is then used as a continuous Cushing stitch to appose visceral pleural surfaces above, f, and, after transfixion at the site of the mattress suture, brought back to the edge of lobe on the inferior surface, g, h, tied and cut, i. A 14 F catheter is placed into the pleural space and the wound closed. After instillation of 100,000 units of penicillin in 5 cc. of saline, catheter is withdrawn.

Arch. Surg. 59:694-704, 1949.



## the new S.K.F. BENZEDREX INHALER

So much better that we have

discontinued 'Benzedrine' Inhaler


Physicians tell us that they and their patients find BENZEDREX INHALER the best inhaler they have ever used.

The active ingredient of Benzedrex Inhaler is 1-cyclohexyl-2-methylaminopropane, a new S.K.F. compound. It has exactly the same agreeable odor as Benzedrine\*, gives even more effective and prolonged shrinkage, and does NOT produce excitation or wakefulness.

We are sure you will find that BENZEDREX INHALER is the best volatile vasoconstrictor you have ever used.

Smith, Kline & French Laboratories, Philadelphia

\*'Benzedrine' (racemic amphetamine, S.K.F.) and 'Benzedrex' T.M. Reg. U.S. Pat. Off,



sponse is improved.

Therapy should be initiated with small, progressively increasing doses of *Decholin Sodium* given intravenously, followed by a course with *Decholin* tablets.

pressure copious, watery bile sluices down the biliary ducts like a spring thaw, carrying off pus, debris, mucus and stagnant bile. With drainage thus re-established the systemic re-

# Decholin

#### brand of dehydrocholic acid

Tablets of 31/4 gr. in bottles of 25, 100, 500, and 1000.

Decholin Sodium® (brand of sodium dehydrocholate) in 20% aqueous solution, ampuls of 3 cc., 5 cc., and 10 cc., boxes of 3 and 20.

DECHOLIN and DECHOLIN SODIUM, trademarks reg. U. S. and Canada.



VITAL STATISTICS

## Maternal Mortality Rates at Record Low

The United States had no more than 1.2 maternal deaths per thousand live births in 1948, a drop of 0.1 from 1947, according to preliminary estimates from state public health agencies. More important than the overall decline in mortality is the fact that the highest state rate in 1948 was 2.7, less than two-thirds of that for the best state in 1933. This figure refutes the charge that rapid improvement in health in this country is limited to the wealthy sections.

J.A.M.A. 141:333-334, 1949.

STATISTICS

c

#### Health of Sailors Improves

The disease rate of the U.S. Navy was lower in 1948 than at any time since records were started in 1850. A major factor in the health record, according to Rear Admiral C. A. Swanson, Navy Surgeon General, was the drop in venereal cases from 85.8 per 1,000 in 1947 to 66.6 in 1948.



PUBLIC HEALTH

#### 9,000 Free Wigs

The British Ministry of Health reports that 4,500 Britons have ordered wigs under the national health plan. Each Briton who obtains a doctor's certificate that he needs a wig is entitled to two.

TREATMENT

#### Xanthine Dust for Asthma

Micropowdered aminophylline or theophylline inhaled in small amounts relieves asthmatic attacks more rapidly than large intravenous doses. From 5 to 60 mg. in a lactose vehicle is taken as needed with a specially designed inhalator. Dr. George V. Taplin and associates of the University of California, Los Angeles, obtained good results in 30 of 35 chronic cases. Slight asthma was checked for one to eight hours and severe spasm alleviated for twenty to forty minutes.


Ann. Allergy 7:513-523, 1949.

PEDIATRICS

#### Moth Balls Cause Anemia

Severe hemolytic reaction may be caused by a reputedly harmless household article, naphthalene moth balls. Drs. Wolf W. Zuelzer and Leonard Apt of Wayne University, Detroit, report 4 cases in which two-year-old children acquired fulminating hemolytic anemia and hemoglobinuria from sucking moth balls. Symptoms appearing in one to four days included listlessness, loss of appetite, diarrhea, vomiting, and pallor. All patients recovered after blood transfusions, sodium bicarbonate, and sodium lactate.

J.A.M.A. 141:185-190, 1949.



The complexities and fast pace of modern living make it all the more essential that mental acuity be sustained throughout the day, including such times as the pre-noon hour when under many conditions mental acuity may wane. Driving high-powered motor cars in congested traffic, working at intricate machines in industry, the pressing demands of business activities, even the demands of household duties, present potential hazards demanding—for safety's sake—that mental acuity be constantly at highest pitch. In children too, alertness must be at maximum intensity at all times to prevent accidents during hours of play.

#### A Good Breakfast - Improved Mental Acuity

In a recent study\* conducted jointly by the departments of physiology and nutrition of a prominent medical college, a direct relationship between late-morning mental acuity and breakfast habits was definitely established. The experiments were planned to learn whether eating breakfast, and eating no breakfast or taking coffee only, made a difference in the subjects' mental acuity during the pre-noon hour.

Six young women graduate students were used as subjects and mental acuity was determined by measurement of simple and choice reaction times to light stimuli. These experiments definitely showed:

ct

rn

at

he

uг

ty

aof

e-

d-

be

0.

11

y.

le

d

l.

r

e

1

- (1) That the *omission* of breakfast increases reaction time, hence acts detrimentally;
- (2) That when breakfast is eaten, reaction time is reduced—hence favorably influenced.

In other words, eating breakfast better prepares one for the demands of living and, from a practical standpoint, serves as a safety precaution against potential accidents due to breakfast skimping or skipping.

#### **Breakfast and Sound Nutrition Planning**

For the foundation of an adequate breakfast, nutrition and health authorities recommend a basic breakfast pattern of fruit or juice, cereal, milk, bread and butter. The cereal serving—breakfast cereal and milk—has many outstanding nutritional advantages. It contributes biologically complete protein; the B vitamins, thiamine, riboflavin and niacin; the minerals, calcium, phosphorus and iron; and needed food energy. It is bland and easily digested and offers many varieties of form, consistency and taste. Its marked economy is a distinctive feature.



 Reprint of the study will be sent on request.



The presence of this seal indicates that all nutritional statements herein have been found acceptable by the Council on Foods and Nutrition of the American Medical Association.

### CEREAL INSTITUTE, INC.

135 South La Salle Street . Chicago 3



A RESEARCH AND EDUCATIONAL ENDEAVOR DEVOTED TO THE BETTERMENT OF NATIONAL NUTRITION

EXPERIMENTAL MEDICINE

#### Treatment of Aneurysms

Periaortic injection of 0.5% dicetyl phosphate in olive oil produces a gradual fibrosis and obliteration of large fusiform aneurysms in animals and might be adapted to the treatment of thoracic and abdominal aneurysms of syphilitic and arteriosclerotic origin. Dr. J. K. Berman and associates of the Indianapolis General Hospital believe that a 0.9% solution is probably the best concentration for treatment of human beings. The mixture is heated and injected beneath the pleura and peritoneum around the thoracic and abdominal aortas. A curved number 18 needle with a blunt bevel is used. The size of the aneurysm determines the curvature of the needle used. A tube is employed to hold the needle steady and avert leaks into pleural or peritoneal cavities. Extravasation of the solution is prevented by manual pressure of firm packing at the site of injection. A small piece of gelfoam is placed over the puncture wound. The dicetyl phosphate solution may also be used to produce vascular constriction for investigation of coronary disease, portal hypertension, renal ischemia, and constricting pericardi-

1. Indiana M.A. 42:889-893, 1949.



ORTHOPEDICS

#### **Strength of Bones**


Human bone is twice as strong as seasoned hickory; one-fourth as strong as cast iron. Unit cubes from the long bones of arms and legs will stand a strain of 23,000 lb. in a compression test before crushing, according to the National Bureau of Standards. Human bone is comparatively weak against twisting strains and lets go at about 3,000 lb. a square inch.

ANTIBIOTICS

#### Polymyxin B for Meningitis

When other measures fail to cure influenzal meningitis, employment of polymyxin B may be lifesaving. Dr. B. M. Kagan of Michael Reese Hospital, Chicago, reports recovery of an infant, thirteen months old, who had had influenzal meningitis four weeks with no sign of improvement despite treatment with streptomycin and sulfadiazine and, finally, with specific rabbit antiinfluenzal serum. These medications were then stopped and polymyxin B was given intramuscularly every four hours in the amount of 50,000 units per kilogram per twenty-four hours. During the first day 1 mg. of polymyxin B was given intrathecally and on the next two days 3.5 mg. was instilled daily. On the fourth day of the new therapy the baby seemed to be more toxic than before, but improvement thereafter was rapid without symptoms or signs or toxicity. Polymyxin B was discontinued after six days. Examination of the child three months after recovery revealed no neurologic or other physical defects.

Pediatrics 4:319-322, 1949.



## Effective Salivary Levels of Tyrothricin

Used as recommended, one Lozille maintains for approximately one-half hour salivary tyrothricin levels as shown in chart.

as

om vill

mrd-

ndely

ets

h.

ire

of

Dr.

OS-

an ad

eks ite ul-

fic ese nd cu-

nt er rst

en

vo

n

he

m

er

ns

nof

ry er

1

The sustained salivary concentrations insure broad antibacterial action against gram-positive organisms responsible for acute oropharyngeal infections. Tyrothricin, unlike topical penicillin, is remarkable for its *lack of local toxicity*. Pleasant-tasting, Lozilles also provide propesin, for non-toxic, long-lasting analgesia.

Each Lozille contains 2 mg. of tyrothricin and 2 mg. of propesin.

Supplied in vials of 15 Lozilles.

LOZILLES

(Lah-Zeels) TYROTHRICIN-PROPESIN LOZENGES

WHITE LABORATORIES, INC., Pharmaceutical Manufacturers, Newark 7, N. J. Free Flow with



# Heparin/ Pitkin Menstruum

'WARNER'

an anticoagulant preparation with prolonged action for the prevention and treatment of thromboembolic disorders.

HEPARIN/PITKIN MENSTRUUM 'Warner' is a safe and clinically established means of providing prolonged anticoagulation action in the body.

One subcutaneous injection of HEPARIN/PITKIN MENSTRUUM 'Warner' is usually sufficient to increase the blood coagulation time for a period of 24 to 48 hours ... without the necessity for the cumbersome, discomforting and time-consuming procedures usually required when maintaining blood fluidity in thromboembolic disease

## Heparin / Pitkin Menstruum



is available in 200-mg and 300-mg ampuls for subcutaneous injection, cartons of 6 ampuls each, with or without vasoconstrictors.

WILLIAM R. WARNER & CO., INC. NEW YORK ST. LOUIS LOS ANGELES GRANTS

## To Keep Young Doctors on Faculty Posts

In an attempt to offset the lure of private practice and uphold teaching standards, the Markle Foundation has granted \$940,000 to medical colleges. Bulk of the funds, states John M. Russell, New York City, will be used to assist 28 young doctors in faculty positions. Institutions benefited include: Bowman Gray School of Medicine, Duke University, New York University, Medical College of Virginia, University of Illinois, Washington University, Yale University, University of Pennsylvania, University of California, and the Medical Film Institute.

GRANTS

#### Rheumatic Fever Research

Seven New York medical colleges and research groups have received grants totaling \$200,000 from the Masonic Foundation for Medical Research and Human Welfare. The funds are to be used to forward study on the cause and cure of rheumatic fever.

BIOCHEMISTRY

#### **Short Cut to Cortisone**

The necessity for using osmium textroxide, a rare and expensive chemical, in the synthesis of cortisone may be eliminated. Scientists of the research staff of Glidden Company, Cleveland, have announced development of a method of creating cortisone from the soybean which may ultimately make the drug relatively plentiful and less expensive than at present.

RESEARCH

#### Institute for Pediatrics

The Playtex Park Research Institute has recently been formed to encourage study of child development and diseases of children. Funds of the organization will be used entirely to augment research in existing institutions. The institute program is supported by the International Latex Corporation, Dover, Del. The governing board, which will administer the finances and supervise all projects, is composed of the following physicians:

Dr. Katherine Bain, U.S. Children's Bureau, Washington, D.C.; Dr. Sidney Farber and Dr. Charles A. Janeway, Children's Medical Center, Boston; Dr. John P. Hubbard and Dr. Joseph Stokes, Jr., Children's Hospital, Philadelphia; Dr. Eva Landsberg, New York City Department of Health; Dr. Milton I. Levine and Dr. Samuel Z. Levine, New York Hospital, New York City: Dr. Charles F. McKhann, Jr., University Hospital of Cleveland; Dr. Henry G. Poncher, University of Illinois, Chicago; Dr. Bret Ratner, Flower and Fifth Avenue Hospital, New York City; Dr. Milton J. E. Senn, Yale University, New Haven, Conn.; Dr. Ashley A. Weech. University of Cincinnati; Dr. Myron E. Wegman, Louisiana University, New Orleans; Dr. James L. Wilson, University of Michigan, Ann Arbor.

PUBLIC HEALTH


#### WHO to Fight Plague

India has been proposed as the first demonstration ground for plague cradication by an international team of health experts from the World Health Organization.

The pioneer investigations in the study of hematology by the eminent Doctor William B. Castle, have contributed immeasurably in the further study and development of medicinals for the treatment of all types of anemias. Doctor Castle's early observation of an existing extrinsic and intrinsic factor assisted greatly in clarifying the normal processes of hemopoiesis. The Armour Laboratories, as a pioneer in the field of endocrinology, is keenly appreciative of Doctor Castle's contributions in the study of hematopathy

Twelfth in the series, PÖRTRAITS OF PIONEERS in Endocrinology. A full-color reproduction of this painting suitable for framing, is available upon request. On your professional letterhead, please address:





## Dr. William B. Castle, 1897-

Dr. Castle was graduated by the Harvard Medical School in 1921. Four years later he began a career in clinical investigation in the field of blood diseases at the Thorndike Memorial Laboratory of the Boston City Hospital. He is director there and Professor of Medicine in Harvard.

Following the great discovery by Minot and Murphy of the beneficial effect of liver feeding, Dr. Castle and his several associates began a series of carefully controlled observations which have demonstrated that the nutritional deficiency in pernicious anemia is "conditioned" by the lack of gastric secretion characteristic of this disease. Their experiments have shown that normal erythropoiesis in man requires both a food (extrinsic) and a gastric (intrinsic) factor and that deficien-


cies of these factors, together with defective intestinal absorption, are the reasons for other macrocytic anemias such as those of sprue and pregnancy. Dr. Castle and his associates have shown that the food (extrinsic) factor is related to Vitamin B<sub>12</sub>.

In Puerto Rico, in 1931, Dr. Castle and his associates were able for the first time to control the macrocytic anemia and diarrhea of tropical sprue by injections of liver extract. Also demonstrated was the striking effectiveness of iron in the treatment of the hypochromic anemia of hookworm disease with or without the removal of the parasites. During the past few years Dr. Castle, in collaboration with others, has conducted experimental studies of the mechanisms of red cell destruction in various types of hemolytic anemias.

NEW DEVICES

#### Radar-like Device Detects Foreign Objects in Body

Ultrasonic energy may be successfully employed to detect and localize calculi or other foreign bodies which are lodged in soft tissues of the body and are not visible by roentgenography. The technic for applying the radar principle to medicine was developed by Dr. George D. Ludwig at the Naval Medical Research Institute, Bethesda, Md. High frequency sound waves generated by a quartz crystal are transmitted into the body tissue from an instrument in direct contact with the skin. Reflections or echoes of these waves occur from the bones and any foreign substance that possesses acoustical properties unlike those of the surrounding tissues. The reflected waves are transformed into electric impulses which are amplified and displayed on a cathode ray oscillograph screen. The distance of the echo from the initial pulse gives the depth of the foreign body in the tissues. Although not yet tried on human beings, the method has been highly successful with animals, and the ultrasonic energy has not been harmful to the tissues.



UROLOGY

#### Treatment of Urethral Stricture

Dilatation of urethral strictures by insertion and retention of progressively larger catheters is an effective method of treatment. Anesthesia is usually unnecessary. Trauma is not as great as with immediate, complete dilatation and the tendency to rapid contracture is lessened. Bleeding and periurethral complications are decreased and the catheter in situ provides opportunity to clear up pyuria by daily irrigations. The disadvantage of prolonged hospitalization is offset by more enduring results. Recently Drs. R. Grant Reid and Claude A. Moore of Montreal General Hospital have gradually dilatated urethras of 7 patients with early results surpassing those obtained by other methods. After local anesthesia, a filiform is passed. The stricture is then dilated with either LeFort's or Phillips' followers to 14-16 F. and a small soft rubber catheter is inserted. Some patients may require anesthesia for this step. At two- to four-day intervals the catheter is replaced with progressively larger sizes. Penicillin and low doses of a sulfonamide are given routinely. In no instance have hemorrhage, hyperpyrexia, or chills been observed.

Canad. M.A.J. 61:278-280, 1949.

EVENTS

#### Marshall Heads Red Cross

Gen. George C. Marshall is president of the American National Red Cross, succeeding Basil O'Connor. Although Mr. O'Connor's term would not have expired until 1951, he resigned effective October 1 because of pressure of personal affairs.



Because "SUDDEN" is a dangerous word in cases of hypertension...it has become almost instinctive with physicians to prescribe Nitranitol. An ideal vaso-dilator, Nitranitol produces gradual reduction of blood pressure in essential hypertension. Nitranitol maintains lowered levels of pressure for prolonged periods. Virtually non-toxic, Nitranitol is safe to use over long periods of time.

# **NITRANITOL®**

For gradual, prolonged, safe vasodilation



When sedation is desired. Nitranitol with Phenobarbital. (½ gr. Phenobarbital combined with ½ gr. mannitol hexanitrate.)

For extra protection against hazards of capillary fragility. Nitranital with Phenobarbital and Ratin. (Combines Rutin 20 mg. with above formula.)

DIAGNOSIS

# Spontaneously Precipitable Protein in Polyarteritis

When protein precipitate appears spontaneously and persists in the serum of a patient who has a bizarre clinical picture, polyarteritis nodosa should be suspected. Dr. Harold Lepow and associates of Lincoln Hospital, New York City, observed the spontaneous appearance of a precipitate at 4° C. in the serum of a patient with proved polyarteritis nodosa. The precipitate differed from previously reported cold fractions in not redissolving at room temperature. The precipitate may also be found in other cases in which the albumin/globulin ratio is reversed, the cephalin flocculation reaction is strongly positive, and serum nitrogen is high.

Am. J. Med. 7:310-316, 1949.

PATHOLOGY

## Platelet Adhesiveness

Use of a braided Fiberglas wick as an adsorbing filter for separation of adhesive from nonadhesive platelets simplifies enumeration of the relative proportions in the total platelet count, and may aid detection of predisposition to thrombosis. Drs. Sylvan E. Moolten and Leo Vroman of St. Peter's General Hospital, New Brunswick, N. J., find that the short exposure of blood on a large surface area tends to reduce the factor of platelet lysis which may cause error in the final calculations. A uniform wait between withdrawal and filtration of the blood is desirable since a longer or shorter interval leads to discrepancies in results.

Am. J. Clin. Path. 19:701-709, 1949.

NEW DEVICES

# **Balloon Tampon Hemostasis**

Bleeding from esophageal varices may be controlled by means of a double balloon tube. The device used by Drs. Thomas B. Patton and Charles G. Johnston of Wayne University, Detroit, consists of 4 ft. of four-lumen plastic tubing. A Rehfuss tip small enough to pass through the nares is affixed to the largest lumen. Separately inflatable balloons are tied on; one 8 in. from the tip and the other, larger, immediately above and contiguous with the first. The third lumen may be used to introduce thrombin just above the balloons. The tube is passed through the nose into the stomach. The position is confirmed by aspiration, the lower balloon inflated to a diameter of 3 in. and pulled snug against the cardia. The tube is then taped to the nose and the upper balloon inflated with 200 to 250 cc. of air. The apparatus is left in place twelve hours with constant suction from the stomach; then the balloons are deflated. If bleeding has stopped, material aspirated from the stomach will be free of blood. The tube is left in place and the patient is given sips of fluid. Nourishment is gradually increased until a full diet is being ingested by the third day. Supportive treatment is necessary.

pro

acc

Arch. Surg. 59:502-506, 1949.



# for Contraception

# LYGENES® LYGEL®

Contraceptive Products provide clinically proved protection

Normal function without anxiety, fear or devices—plus patients

The aim of modern contraception—to instill complete comfidence—is fully realized in this line of contraceptives which provide a high degree of effectiveness. Non oxic, esthetically acceptable, non-irritating, economical.



ecial Formula Corporation
DISTRIBUTOR

Park Avenue, New York 22, N.Y.

LYGENES VAGINAL SUPPOSITORIES Clinically Proved Highly Effective. Small, nonodorous vaginal suppositories which form an adhesive, effective cervical barrier in a matter of minutes. No diaphragm or other devices required. Convenient. Facilitate patient-cooperation. Economical—in boxes of 12, foil-wrapped.

ACTIVE INGREDIENTS

Hydroxyquinoline Benzoate 0.30% p-Chloro-symm.-m-dimethylhydroxybenzene 0.05% p-tert. Amythydroxybenzene 0.05% Zinc Sulfocarbolate 0.50% pH 4 (when dispersed in 4 parts normal saline)

LYGEL VAGINAL JELLY—A jelly of high spermicidal efficacy with freedom from irritation and ready patient-acceptance. Readily dispersible. Does not lose viscosity at body temperature. 3-oz. tubes.

### ACTIVE INGREDIENTS

p-Chloro-symm,-m-dimethylhydroxybenzene 0.05% p-tert. Amylhydroxybenzene 0.05% Benzalkonium Chloride 0.10% Lactic Acid 0.25% p.H. 3.4

Literature and clinical trial packages on request.

Special Formula Corporation, De 445 Park Avenue, New York 22, You may send me (check your pr	N. Y.	
1 Package LYGENES Sup		
1 LYGEL Refill		
		M.D.
ADDRESS.		
CITYZONE	STATE	

# Washington Letter

(Continued from page 41)

the last days of Congress, the point has not been resolved.

The American Medical Association and some other organizations are strongly opposed to the principle of free treatment without a means test.

However, sentiment in Congress is overwhelmingly in favor of a school health bill. This bill passed the Senate without debate. It was tied up in a House committee, mostly because of the "means" issue. If it is not the law of the land now, it is certain to be shortly after the first of the year when the second session meets.

Under this "no means-test" clause, every private physician would be assured payment for treatment of school-age children. For a few years, some states might be expected to require an ability-to-pay test, but political pressure would bring all states into the "free treatment" class in a few years. It would, automatically, bring many millions of patients under a form of federal health insurance.

5 The direction this program takes will be largely determined by the private physician and professional associations. The law defines a specific role for the private physician in treatment of patients and for medical associations in administration.

The school health plan is the closest approach yet to a federal health program. It offers the physician an opportunity to observe the workings of the plan and to take a hand in formulation of policies.



"I'm sick of normal deliveries. This time I want a cesarean."



# Iron in adequate dosage

"is almost a physiological necessity in infancy and childhood and [childbearing] women . . ."

Sundaram, S.K.: Lancet. 1:568, 1948



# and Feosol is iron in its most effective form

Feosol—both the Tablets and the highly palatable Elixir—contains adequate dosage of ferrous sulfate, grain for grain the most effective form of iron. Feosol is easily absorbed and readily tolerated. Feosol effects rapid hemoglobin regeneration and prompt reticulocyte response.

Smith, Kline & French Laboratories, Philadelphia



# Feosol Tablets Feosol Elixir

the standard forms of iron therapy

# Million Volt Cancer Therapy

Veterans Administration shortly will have ready the results of a continuing study of veterans who received million-volt x-ray treatment for cancer. Three years ago about 150 men received the treatments on an experimental basis. VA rounded up as many of them as possible for post-treatment studies, to determine the long-time benefits or harms of this type of therapy. The study is expected to throw some light on just how much x-ray the body can take for maximum benefit and minimum harm. The investigation is under direction of Dr. Aubrey O. Hampton of Garfield Hospital, Washington, D.C.

# Appeal to Druggists

Federal Security Administrator Oscar Ewing, addressing the National Association of Retail Druggists, made a direct dollars-and-cents argument to win support for national health insurance. He said in part:

Certainly, every gain in our public health and other services is a gain for the drug trade. For the drug business flourishes most in those areas where the concern for health is the greatest. And the partnership of the druggists with the local public health doctor is as real as with the private physician.

For the last month of Congress, almost nothing was heard of the President's health plan. However, supporting studies are being revised in preparation for the next session.



# With Nose and Throat Medications Prescribe DeVilbiss Atomizers Better Penetration—



)5-

al de nt

th

lic

ess he ad th

r,

WHEN home treatment of nose and throat infections is indicated, DeVilbiss Atomizers offer your patients the most effective means of therapy.

X-Ray research confirms that a sprayed solution has superior penetration and wider distribution. The result, proved by clinical investigation, 1 is a two-fold longer period of relief.

DeVilbiss Atomizers may be prescribed with complete confidence based on the sixty-one years DeVilbiss has served the medical profession.



# Reference:

1. PERSKY, A. H.:
"Penicillin-Vasoconstrictor Treatment of Post-Influenzal Rhinitis
and Sequellae",
Medical Record,
November, 1947.



THE DEVILBISS COMPANY
Toledo 1, Ohio

# Welfare Plan Redrafted

President Truman, turned down once on his plan for a cabinet department of education, health, and welfare, is trying again. The plan is being redrafted at the White House for presentation to Congress after the first of the year.

It is understood that some arrangement is being made to placate the private welfare organizations which felt that their activities would suffer under the proposed plan. However, the administration will not back down on including health activities in the new department. Last session this brought it into head-on conflict with American Medical Association and other professional organizations.

# Atomic Research Accelerated

Russia's possession of the atomic bomb is having one valuable reaction: It has insured that basic atomic research will not be slighted for years to come. Funds for atomic research were trimmed slightly by the last Congress. However, a week after announcement that Russia had the bomb, a move was underway to get more money for basic research, including medical research. Sen. Brian McMahon, chairman of the joint atomic congressional committee, joined scientists in this appeal: We have made use of all the basic knowledge we had three years ago. In some fundamental phases of the subject our ignorance is abysmal.



### No. 9863 Steeltone Table

This large, efficient table has 28 different features that save you time and effort.

# Give Your Office The "New Look" Too

Now, with the new Hamilton Steeltone Suite, you can modernize your office and make it more inviting to patients. Modern in style, with superior Hamilton all-welded construction, this new furniture says "Welcome" to patients in a friendly and dignified way. Send in the coupon for more information.

# Hamilton Manufacturing Co. Two Rivers • Wisconsin

This coupon will bring you a copy of Hamilton's new catalog showing three modern office suites. Send it in today.

Send me information on the St	teeltone as soon as available.	
		M.D.
Address		
City & State		MM-11-49

# THROAT SPECIALISTS REPORT ON 30-DAY TEST OF CAMEL SMOKERS—

# "Not one single case of throat irritation due to smoking CAMELS!"



YES, these were the findings in A total of 2,470 weekly examinations of hundreds of men and women from coast-to-coast who smoked only Camels for 30 consecutive days! And the smokers in this test averaged one to two packages of Camels a day!



According to a Nationwide survey:

# MORE DOCTORS SMOKE CAMELS

than any other cigarette!

When three leading independent research organizations asked 113,597 doctors what eigarette they smoked, the brand named most was Camel!

R. J. Reynolds Tobacco Co., Winston-Salem, N. C.

# Current Books & Pamphlets

This catalogue is compiled from all available sources, American and foreign, to insure a complete listing of the month's releases.

### Medicine

THE DIAGNOSIS OF PANCREATIC DISEASE by Louis Baumann. 74 pp., ill. J. B. Lippincott Co., Philadelphia. \$5

CLINICAL METHODS by Sir Robert Hutchison and Donald Hunter. 12th ed. 488 pp., ill. Cassell & Co., London. 17s. 6d.

HANDBOOK OF DIGESTIVE DISEASES by John Leonard Kantor and Anthony M. Kasich. 2d ed. 658 pp., ill. C. V. Mosby Co., St. Louis. \$11

THE VALUE OF HORMONES IN GENERAL PRACTICE by William N. Kemp. 115 pp. Burgess Publishing Co., Minneapolis. \$3

STOMACH DISEASE AS DIAGNOSED BY GASTRO-SCOPY by Eddy D. Palmer. 200 pp., ill. Lea & Febiger, Philadelphia. \$8.50

BEDSIDE DIAGNOSIS by Charles MacKay Seward. 376 pp., ill. E. & S. Livingstone, Edinburgh. 17s. 6d.

### Surgery

ADVANCES IN SURGERY, VOLUME 1 edited by William DeWitt Andrus et al. 566 pp., ill. Interscience Publishers, New York City. \$11

BLOOD TRANSFUSION edited by Geoffrey L. Keynes. 574 pp., ill. Williams & Wilkins Co., Baltimore. \$12.50

# 93% PROTECTION<sup>1</sup>

IN BRONCHIAL ASTHMA



\* An especially purified synthetic racemic epinephrine available with Vaponefrin nebulizer, hand bulb, and carrying case (VAPONEF-RIN AEROSOL UNIT).



# **VAPONEFRIN® SOLUTION**

Of all the common bronchodilators, Vaponefrin Solution\* provides the greatest degree of protection for the longest period of time against histamine-induced bronchospasm.

1. Segal, M. S.: Dis. Chest 14: 795-823, 1948.

# VAPONEFRIN COMPANY

6816 MARKET STREET UPPER DARBY, PENNSYLVANIA

# Write FOR



ANGINA PECTORIS

# THESODATE Grewer

Theobromine Sodium Acetate (71/2 gr.) with Phenobarbital (1/4 gr.) enteric coated for improving cardiac action and coronary circulation. Other strengths and combinations also available.

LUETIC HEART



Potassium Iodide (15 gr.) enteric coated preferable to the solution wherever potassium iodide is indicated including its use in tertiary syphilis.

CARDIAC **EDEMA** 




Ammonium Chloride (15 gr.) enteric coated for the treatment of cardiac edema with half the usual number of tablets.

Literature and Samples on request.



COMPANY, WORCESTER 8, MASSACHUSETTS U.S.A.



Yopa-Cresolene is recommended in Bronshirls, Bronchial Asthma, Spasmodic Croup, Whooping Cough. Excellent for children's stuffy nasol colds.

PRESCRIBED SINCE 1870
Send for special brochure

THE VAPO-CRESOLENE CO.
62 Cortlandt St. New York 7, N. Y.

# LITTLE TOIDEY





# for Training the Baby

Illustrated are Little Toidey on adult seat (also fits on low base) \$5.50 retail, and Toddler's Toidey (base and pan) \$3.95 retail.

20% COURTESY DISCOUNT from regular retail price (and prepaid in U.S.A.) to physicians for office or home on any or all of the items in the complete Toidey Training Unit: Little Toidey (wood or plastic) Toidey Base, Toideyette (deflector) Specimen Collector, Two-Steps. Write for complete list and free book "Training the Baby."

The TOIDEY COMPANY

Gertrude A. Muller, Inc.

FORT WAYNE • INDIANA

DAS AKUTE ABDOMAN: KLINIK UND THER-APIE DER AKUTEN LEBENSBEDROHENDEN BAUCHERKRANKUNGEN by Hubert Kunz. 357 pp., ill. Urban & Schwarzenberg, Vienna. 105 Sch.

TEXT BOOK OF SURGERY by Patrick Kiely. 1,184 pp., ill. H. K. Lewis, London.

455.

## Allergy

PROGRESS IN ALLERGY, VOLUME II edited by Paul Kallós. 356 pp., ill. Interscience Publishers, New York City. \$7.50

PRÉCIS DES MALADIES ALLERGIQUES by Pasteur Vallery-Radot. 223 pp., ill. Editions Médicales Flammarion, Paris. 550 fr.

### Anatomy

some notes on galen's anatomy by Wynfrid Laurence Henry Duckworth. 42 pp. W. Heffer & Sons, Cambridge. 2s. 6d.

ANATOMY OF THE FEMALE PELVIS by Frederick Arthur Maguire. 4th ed. 176 pp., ill. Angus & Robertson, Sydney. 42s.

VERTEBRATE BODY by Alfred Sherwood Romer. 643 pp., ill. W. B. Saunders Co., Philadelphia. \$5.50

### Anesthesia

GAS AND AIR ANALGESIA by R. J. Minnitt. 4th ed. 86 pp., ill. Baillière, Tindal & Cox, London. 5s.

# Gynecology & Obstetrics

HAVING YOUR BABY: MODERN INSTRUCTIONS FOR EXPECTANT MOTHERS by Leonard H. Biskind. 96 pp. Western Journal of Surgery Publishing Co., Portland, Oregon. \$2.50

THE MIDWIFE'S TEXT-BOOK by R. W. Johnstone, 4th ed. 400 pp., ill. A. & C. Black,

London. 20s.

BIRTH CONTROL TODAY by Marie C. Stopes. 9th ed. 242 pp., ill. Alexander Moring, London. 6s.

### Proctology

TREATMENT IN PROCTOLOGY by Robert Turell. 248 pp., ill. Williams & Wilkins Co., Baltimore. \$7

### **Pharmacology**

LECTURE NOTES ON PHARMACOLOGY by Joshua Harold Burn. 128 pp. Charles C Thomas, Springfield, Ill. \$1.60

(Continued on page 158)

Complete Stability . . .

Outstanding Palatability . . .

# Syrup Choline Dihydrogen Citrate



(Flint)

Continued patient acceptance of choline over prolonged periods can be assured by prescribing—



# "Syrup Choline" (Flint)

25 per cent W/V—containing one gram of choline dihydrogen citrate in each 4 cc.

Supplied in pints and gallons.

# "Capsules Choline" (Flint)


containing 0.5 gram of choline dihydrogen citrate. Supplied in bottles of 100, 500 and 1000.

"Choline (Flint)" is indicated in fatty infiltrations of the liver associated with alcoholism, infectious hepatitis, early cirrhosis, diabetes, malnutrition.

For your copy of "The Present Status of Choline Therapy in Liver Dysfunction"—write



# NOW...ALL OF 1949's NEW ONE HUGE, ILLUSTRATED



- Over 800 Informative Pages
- 420 Abstracts by 700 Authors
- 24 Interesting Diagnostix
- 45 Special Articles & Exhibits
- Hundreds of Fine Illustrations
- Beautifully Printed Richly Bound
- Double Indexed By Author and By Subject

# SPECIAL Send No

ach year, for the past seven years, we have published the MODERN MEDICINE ANNUAL—the one single volume that keeps doctors "up-to-date" on the year's medical progress! During that time, thousands of physicians and surgeons have come to regard the Modern Medicine Annual as an indispensable reference to contemporary medical thought . . . a "most valued assistant" in keeping abreast of medicine's fast pace!

# PUBLISHED SOON-LIMITED EDITION OF THE 1950 ANNUAL

Now-we are planning to publish a limited edition of the Modern Medicine Annual for 1950-which will be ready for distribution shortly after January 1st. As in the former editions, the 1950 Modern Medicine Annual contains hundreds of authoritative articles on new surgical techniques, new instruments and equipment, new medicines, new theories and methods that were developed during the year, In its 800 informative pages, you will find all of the important reports, digests and abstracts that ap-

MODERN

# MEDICAL DISCOVERIES IN FULLY-INDEXED VOLUME

# **ODERN MEDICINE ANNUAL**

Alire-Publication Price . . . \$500

No Money — Just Mail the Reservation Card

peared in "Modern Medicine" during 1949. Each article is clear, concise and easy to understand-and there are hundreds of illustrations to emphasize and complement the text!

even

hed

INE

ngle

tors

ear's

that

cians

o re-

cine

efer-

dical

lued

st of

DF

pub-

the 1 for

uary

ions,

icine

is of

new

stru-

new

and

oped

0 infind

orts.

ap-

# MANY SPECIAL FEATURES

In addition to the 420 abstracts-there are 145 special articles and exhibits on the very latest discoveries in the various fields of medicine-plus 24 interesting "Diagnostix." Furthermore, this valuable book is so completely indexed that you can locate the information you want in a matter of seconds! One 16-page index lists all of the authors and another 40-page index covers the entire book by subjects! Truly, all of 1949's medical progress . . . at your fingertips!

# A WORTHY ADDITION TO YOUR LIBRARY

Beautifully printed and richly bound, the y for 1950 Modern Medicine Annual is a handsome—as well as a useful—addition to your medical library. Many physicians order two copies-one for leisure reading at home, and one for the office for quick reference! That way, you are always sure of finding the information you want, whenever you want it!

# SEND NO MONEY-HIST THE RESERVATION CARD

Yes, the 1950 Modern Medicine Annual is literally packed with authoritative medical information-facts in diagnosis and treat-

ment that will help you every day in your practice! We have already received thousands of advance orders from doctors who have purchased previous issues-and who know what an invaluable reference work it is! Since only a limited edition of this 1950 Annual will be printed—we urge you to reserve your copy NOW. Send no moneyjust mail the postage-paid Reservation Card . . . and you will receive your copy shortly after January 1st-at the special pre-publication price of only \$5.00 (instead of the regular price of \$6.00)! Remember-this special pre-publication offer is necessarily limited-so be sure to mail the Reservation Card today!


# 10 DAYS FREE TRIAL EXAMINATION

So confident are we that you will find this 1950 Modern Medicine Annual an indispensable guide in your everyday practice that we make this unusual offer. Send no money-just the Reservation Card. When your copy arrives (shortly after January 1st), read it at your leisure. Note what a helpful, reliable, time-saving reference work it is. Then, if you do not agree that this is the easiest method of keeping "upto-date" on the year's medical progressreturn your copy within 10 days and your money will be promptly refunded!

> Don't delay! The edition is limited so mail the Reservation Card NOW.

NMEDICINE 510 ESSEX BLDG., MINNEAPOLIS 3, MINN.

# NOW...ALL OF 1949's NEW ONE HUGE, ILLUSTRATED



- Over 800 Informative Pages
- 420 Abstracts by 700 Authors
- 24 Interesting Diagnostix
- 45 Special Articles & Exhibits
- Hundreds of Fine Illustrations
- Beautifully Printed Richly Bound
- Double Indexed By Author and By Subject

# THE 1950 SPECIAL

Send No

Each year, for the past seven years, we have published the MODERN MEDICINE ANNUAL—the one single volume that keeps doctors "up-to-date" on the year's medical progress! During that time, thousands of physicians and surgeons have come to regard the Modern Medicine Annual as an indispensable reference to contemporary medical thought . . a "most valued assistant" in keeping abreast of medicine's fast pace!

# PUBLISHED SOON— LIMITED EDITION OF THE 1950 ANNUAL

Now-we are planning to publish a limited edition of the Modern Medicine Annual for 1950-which will be ready for distribution shortly after January 1st. As in the former editions, the 1950 Modern Medicine Annual contains hundreds of authoritative articles on new surgical techniques, new instruments and equipment, new medicines, new theories and methods that were developed during the year, In its 800 informative pages, you will find all of the important reports, digests and abstracts that ap-

MODERN

# MEDICAL DISCOVERIES IN LY-INDEXED VOLUME

# RN MEDICINE ANNUAL E-PUBLICATION PRICE . . .

Just Mail the Reservation Card Send No Money - Just This Reservation Card

MODERN MEDICINE ANNUAL 84 South 10th St., Minneapolis 3, Minn.

Date

 Please reserve for me a copy of the 1950 Modern Medicine Annual. It is understood that I will be billed for this at the special pre-publication price of only \$5.00, plus a few cents postage and delivery charges. It is further understood that if I am not completely satisfied with my copy, I may return it within 10 days for prompt refund.

Name

Please print name

an Address

Vo

even hed NE

agle

tors ar's that

ians

o recine

efer-

dical

lued st of

) F

pub-

the 1 for

y for

uary

ions,

icine

is of

stru-

new

and

oped

0 infind

orts,

t ap-

O

City

 Check here if you enclose \$5.00 in an envelope with this Reservation Card.
 We will send you the 1950 Modern Medicine Annual, postpaid. Same money-back guarantee.

Modern Medicine Amium is a manusome—as well as a useful—addition to your medical library. Many physicians order two copies-one for leisure reading at home, and one for the office for quick reference! That way, you are always sure of finding the information you want, whenever you want it! new

# SEND NO MONEY-HUST THE RESERVATION CARD

Yes, the 1950 Modern Medicine Annual is literally packed with authoritative medical information-facts in diagnosis and treatmake this unusual offer. Send no money-just the Reservation Card. When your copy arrives (shortly after January 1st), read it at your leisure. Note what a helpful, reliable, time-saving reference work it is. Then, if you do not agree that this is the easiest method of keeping "upto-date" on the year's medical progressreturn your copy within 10 days and your money will be promptly refunded!

> Don't delay! The edition is limited so mail the Reservation Card NOW.

EDICINE 510 ESSEX BLDG., MINNEAPOLIS 3, MINN.

# NOW...ALL OF 1949's NEW ONE HUGE, ILLUSTRATED



# **THE 1950**

CPFCIAL


ed

FIRST CLASS Permit No. 2139 (Sec. 34.9, P. L. & R.) Minneapolis, Minn.

BUSINESS REPLY CARD
No Postage Stamp Necessary if Mailed in the United States

POSTAGE WILL BE PAID BY-

Modern Medicine Publications 84 So. 10th Street Minneapolis 3, Minn.



- Over 800 Informative Pages
- 420 Abstracts by 700 Authors
- 24 Interesting Diagnostix
- 45 Special Articles & Exhibits
- Hundreds of Fine Illustrations
- Beautifully Printed Richly Bound
- Double Indexed By Author and By Subject

INE ITAU MINIO

Now-we are planning to publish a limited edition of the Modern Medicine Annual for 1950-which will be ready for distribution shortly after January 1st. As in the former editions, the 1950 Modern Medicine Annual contains hundreds of authoritative articles on new surgical techniques, new instruments and equipment, new medicines, new theories and methods that were developed during the year, In its 800 informative pages, you will find all of the important reports, lit digests and abstracts that ap- in

MODERN

# MMEDICAL DISCOVERIES IN FULLY-INDEXED VOLUME

# **NODERN MEDICINE ANNUAL**

AIRE-PUBLICATION PRICE ... \$500

Noney—Just Mail the Reservation Card

peared in "Modern Medicine" during 1949. Each article is clear, concise and easy to understand-and there are hundreds of illustrations to emphasize and complement the text!

# MANY SPECIAL FEATURES

In addition to the 420 abstracts-there are 45 special articles and exhibits on the very latest discoveries in the various fields of medicine-plus 24 interesting "Diagnostix." Furthermore, this valuable book is so completely indexed that you can locate the information you want in a matter of seconds! One 16-page index lists all of the authors and another 40-page index covers the entire book by subjects! Truly, all of 1949's medical progress . . . at your fingertips!

# A WORTHY ADDITION TO YOUR LIBRARY

pub-

the

1 for

y for

uary

ions,

cine

is of

stru-

new

and

oped

0 in-

Beautifully printed and richly bound, the 1950 Modern Medicine Annual is a handsome—as well as a useful—addition to your medical library. Many physicians order two copies—one for leisure reading at home, and one for the office for quick reference! Thatway, you are always sure of finding the information you want, whenever you want it! new

# SEND NO MONEY-HIST THE RESERVATION CARD

find Yes, the 1950 Modern Medicine Annual is orts, literally packed with authoritative medical ap-information-facts in diagnosis and treatment that will help you every day in your practice! We have already received thousands of advance orders from doctors who have purchased previous issues-and who know what an invaluable reference work it is! Since only a limited edition of this 1950 Annual will be printed—we urge you to re-serve your copy NOW. Send no money just mail the postage-paid Reservation Card . and you will receive your copy shortly after January 1st-at the special pre-publication price of only \$5.00 (instead of the regular price of \$6.00)! Remember-this special pre-publication offer is necessarily limited-so be sure to mail the Reservation Card today!

# 10 DAYS FREE TRIAL EXAMINATION

So confident are we that you will find this 1950 Modern Medicine Annual an Indispensable guide in your everyday practice that we make this unusual offer. Send no money-just the Reservation Card. When your copy arrives (shortly after January 1st), read it at your leisure. Note what a helpful, reliable, time-saving reference work it is. Then, if you do not agree that this is the easiest method of keeping "upto-date" on the year's medical progressreturn your copy within 10 days and your money will be promptly refunded!

> Don't delay! The edition is limited so mail the Reservation Card NOW.

N MEDICINE 510 ESSEX BLDG., MINNEAPOLIS 3, MINN.

# Greater COMFORT for the BILIARY Patient

TOROCOL Tablets speed relief from biliary constipation, after-meal fullness, dyspepsia, eructation... by improving bile flow, tolerance to fatty foods, bowel regularity. Successfully prescribed for over 39 years.

# TOROCOL



CHOLERETIC
GENTLE LAXATIVE
FOR THE STAGNANT
GALLBLADDER

For Samples and Literature Write

# THE PAUL PLESSNER COMPANY

DETROIT 26, MICHIGAN



# PARAVOX

VERI-small Hearing Aid, although tiny in size, can produce powerful volume, if needed, for extreme hearing loss. Light in weight, easy to conceal.

Another point, Paravox nation-wide sales outlets permit "one-minute" service, a vital point to users. Thousands enjoy their Paravox, like the exclusive plastic chassis which has been 600 foot drop-tested, ton pressure tested, humidity tested, and is a national design winner.

Accepted by the Council on Physical Medicine, American Medical Association.

Made by PARAVOX, INC. 2056 E. Fourth Street, Cleveland 15, Ohio

### Otolaryngology

FUNDAMENTALS OF OTOLARYNGOLOGY: TEXTBOOK OF EAR, NOSE AND THROAT DISEASES by Lawrence R. Boies. 443 pp., ill. W. B. Saunders Co., Philadelphia. \$6.50

KURZGEFASSTES LEHRBUCH DER ERKRAN-KUNGEN DES OHRES, DER NASE UND DES HALSES by Alfred Güttich. 257 pp., ill. George Thieme, Stuttgart. 16 M.

OTO-RHINO-LARYNGOLOGIE IM KINDESAL-TER: EINSCHLIESSLICH DER ENDOSKOPIE by Elemer Josef Jents et al. 326 pp., ill. Wilhelm Maudrich, Vienna. 96 Sch.

RECENT ADVANCES IN OTO-LARYNGOLOGY by Robert Scott Stevenson. 2d ed. 396 pp., ill. Blakiston Co., Philadelphia. \$6

### Orthopedics

1.E ENERVAZIONI ARTICOLARI by F. Morin and F. Roasenda. 138 pp. Minerva Medica, Turin, Italy. 1400 lire.

FRAKTUREN UND LUXATIONEN by Arthur Schäfer. 394 pp., ill. Wissenschaftliche Verlagsgesellschaft, Stuttgart. 24 M.

A PRACTICE OF ORTHOPAEDIC SURGERY by T. P. McMurray, 3d ed. 444 pp., ill. Edward Arnold & Co., London, 3os.

# Industrial Medicine

INDUSTRIAL TOXICOLOGY by Lawrence T. Fairhall. 483 pp. Williams & Wilkins Co., Baltimore. \$6

INDUSTRIAL TOXICOLOGY by Alice Hamilton and Harriet L. Hardy. 2d ed. 574 pp. Paul B. Hoeber, New York City. \$6.50

### Neurology

ÜBER ZWISCHENHIRNSYNDROME by F. Laubenthal. 52 pp., ill. Georg Thieme, Stuttgart. 3.60 M.

TRAITÉ DE MÉDECINE, TOME XV. MALADIES DU SYSTÈME NERVEUX edited by A. Lemierre et al. 1,254 pp., ill. Masson & Co., Paris. 3400 fr.

TRAITÉ DE MÉDECINE, TOME XVI, MALADIES DER SYSTÈME NERVEUX edited by A. Lemierre et al. 1,251 pp., ill. Masson & Co., Paris. 3,400 fr.

PROGRESS IN NEURO: OGY AND PSYCHIATRY, VOLUME IV edited by E. A. Spiegel et al. 592 pp. Grune & Stratton, New York City. \$10

DISEASES OF THE NERVOUS SYSTEM by Francis M. R. Walshe. 6th ed. 376 pp., ill. E. & S. Livingstone, Edinburgh. 17s. 6d.

# so comfortable

Both physically and psychologically, TAMPAX tampons are amazingly comfortable intravaginal menstrual guards. They cannot induce odor, perineal irritation or infection via rectum. And, with the individualization and convenience of protection provided by the three absorbencies (Regular, Super, Junior), their use is said to tend to make women "forget they are menstruating."\* These dainty cotton tampons are also thoroughly safe

and adequate.

\*West. J. Surg., Obstet. & Gynec., 51:50, 1943; J.A.M.A., 128:490, 1945.

TAMPAX INCORPORATED PALMER, MASS.

the internal menstrual guard



MM-15-119

# NO OTHER RUB GIVES FASTER RELIEF IN RHEUMATIC ACUES-PAINS

# **Lumbago and Neuritis Discomfort**

Musterole is an excellent analgesic, decongestive, and counter-irritant for relieving muscular aches, pains, soreness and stiffness—for helping to break up topical congestion. It has all the advantages of a mustard plaster yet eliminates the fuss and bother of making one, and is far more comfortable for the patient. In 3 Strengths: Children's Mild. Also Regular and Extra Strong for adults.





# Child Psychology

THERAPEUTIC GROUP WORK WITH CHILDREN by Gisela Konopka, 134 pp., ill. University of Minnesota Press, Minneapolis. \$2.50

UNDERSTANDING YOUNG CHILDREN by Dorothy W. Baruch. 51 pp., ill. Columbia University, New York City. \$.60

## **Embryology**

LEHRBUCH DER EMBRYOLOGIE by Walter Brandt. 648 pp., ill. S. Karger, Basel, Switzerland. 56 Sw. fr.

FUNDAMENTALS OF COMPARATIVE EMBRY-OLOGY OF THE VERTEBRATES by Alfred F. Huettner. 309 pp., ill. Macmillan Co., New York City. \$5

# Biography

explorer of the human brain: the life of santiago ramon y cajal by Dorothy F. Cannon. 303 pp., ill. Henry Schuman, New York City. \$4

sir William cowers, 1845-1915 by Macdonald Critchley. 118 pp., William Heinemann Medical Books, London. 178. 6d.

### **Dictionaries**

DIZIONARIO INGLESE-ITALIANO PER LE SCIENZE MEDICHE by Ruggero Marconi and Elena Zino. 565 pp. Edizioni Minerva Medica, Turin. 2400 lire

THE ORIGIN OF MEDICAL TERMS by Henry Alan Skinner. 379 pp. Williams & Wilkins Co., Baltimore. \$7

# Nursing

CLINICAL INSTRUCTION by Amy Frances Brown. 571 pp., ill. W. B. Saunders Co., Philadelphia. \$5.50

WARD ADMINISTRATION AND CLINICAL TEACHING by Florence M. Gipe and Gladys Sellew. 357 pp., ill. C. V. Mosby Co., St. Louis. \$4.25

GYNECOLOGY AND GYNECOLOGIC NURSING by Norman F. Miller et al. 2d ed. 485 pp., ill. W. B. Saunders Co., Philadelphia. \$4.25

MEDICAL AND NURSING DICTIONARY AND EN-CYCLOPAEDIA by Evelyn Clare Pearce. 9th ed. 736 pp. Faber & Faber, London. 17s. 6d.

ward administration by Margaret Randall. 326 pp., ill. W. B. Saunders Co.. Philadelphia. \$4

# at the very first sign of a c

# to development-

antihistaminic therapy has been reported to abort the development of the common cold in 90% of the patients commencing therapy within the first hour of the appearance of symptoms.

# DETOUR distressing symptoms-

antihistaminic therapy shortens the duration and decreases the severity of an established cold.1.3

# spread of infection to others—

the elimination of sneezing, lacrimation, rhinorrhea and coughing reduces cross-infection.'

# RICIT

(Antipyretic-analgesic-antihistaminic)

combines the classical "A.P.C. formula" (Acetylsalicyclic acid 3.5 gr., Acetophenetidin 2.5 gr. and Caffeine 0.5 gr.) with Chlor-Trimeton\* the antihistaminic with minimal side-effects and greater effectiveness in doses as low as 2-4 mg.

The Allergic Concept of the Common Cold: The symptoms of upper respiratory infections closely resemble those found in vasomotor rhinitis and hay fever. More histamine-like substances were found in the nasal secretions of persons suffering from colds than in allergic rhinitis.4

Dosage and Timing: Two Coricidin Tablets at the very first indication of a cold, then one tablet every three or four hours for three or four days. In established colds, one tablet every three or four hours for palliative effect.

Packaging: Coricidin Tablets, tubes of 12, bottles of 100 and 1000.

Bibliography: 1. Brewster, J. M.: Indust. Med. 18:217, 1949. 2. Murray, H. C.: Indust. d. 18:215, 1949. 3. Tislow, R. and others: Federation Proc., Part 1, 8:338, 1949. 4. Troescher-Elam, E.; Ancona, G. R., and Kerr, W. J.: Am. J. Physiol. 145:711, 1945.

\* T. M. Schering Corporation

CORPORATI







SIMPLE - ACCURATE - COMPACT La Mar Cancer Detection Stain Set (for Papanicolaou Method)

123 W. 64th St., New York 23, N. Y.

Complete staining in 10 Minutes

ŀ

CA

L

S

Order through your Surgical Supply Dealer
LA MAR LABORATORIES INC.
24 East 21 Street New York 10. N. Y.

La Mar DIAGNOSTIC SETS



# **PATIENTS**

# I Have Met

The editors will pay \$1 for each story published. No contribution will be returned. Send your experiences to the Patients I Have Met Editor, MODERN MEDICINE, 84 South Tenth St., Minneapolis 3, Minn.

### **Proportional Representation**

A routine history was being taken from a young woman in the gynecology clinic. She was asked if she had noticed anything unusual about her menstrual periods recently.

After some thought, she replied, "No, except that in February my period lasted only two days instead of the ordinary six. I didn't think that unusual because February is a short month."—I.J.M.

### An Opportunist

"I'm so discouraged," sighed the pretty nurse, "Everything I do seems to be wrong." With a gleam in his eye the intern asked, "How about a date tonight?"—s.f.w.

### A Good Dog

I was caught in a heavy storm in the Ozark foothills and had to impose on the hospitality of one of my patients. At dinner I asked for cream for my coffee.

"Haven't had any milk since our dog died," drawled the hillbilly. Then, much to my relief, he added, "A good dog. He always brought in the cows."—s.s.

e t

ree

our

B.

DS

ire-

OW


7hit

B.

ran



"Now exhale."



# B. F. Goodrich: the long and the short of surgeons' gloves

for convenience, choose in a me you know for proction. B. F. Goodrich makes tree types of gloves to meet our requirements:

he on ts.

ny og ch

g.

S.S.

B. F. Goodrich "Miller" rand surgeons' gloves—issue thinness to the finger ps assures sensitive touch, are hand freedom. Full backs, aug fitting wrists to fit around own cuffs. Sizes 6 to 10. 7hite or brown. "Smooth" or Cutinized" surface.

B. F. Goodrich "Miller" rand examination gloves tme touch sensitivity, same protection as "Miller" brand surgeons' gloves. Shorter length cuff. Sizes 7 to 9. White only.

The new B. F. Goodrich "Special Purpose" gloves— Specially created for those who develop an allergic dermatitis when using ordinary rubber gloves. Impressive record of successful use. Tissue thin. Strong. Sizes 6½ to 9½. Look for identifying green band on the cuff.

Whichever type you use, whenever you use them you'll be assured of dependable strength because B. F. Goodrich surgeons' gloves are made by the patented Anode process—a single strong, tissue-thin layer of pure rubber made on sculptured hand forms. Better fit too—and maximum touch sensitivity, greater comfort, plus the true economy of long service. Order B. F. Goodrich gloves from your hospital or surgical supply dealer. The B. F. Goodrich Company, Sundries Div., Akron, Ohio.

B.F. Goodrich
Surgeons' Gloves

# CHLORTUSSIN (ANGIER)

a valuable adjunct treatment for coughs due to common cold, bronchitis, pharyngitis, laryngitis, pertussis. Especially designed for children. Contains B<sub>1</sub>. Bottles of 4 and 8 fl. ozs.

# ANGIER CHEMICAL CO., BOSTON 34, MASS. Makers of

THI-BO-MIN — pleasant, non-alcoholic iron-liver preparation plus B<sub>1</sub> for supplemental treatment of *Deficiency Anemias*. Suitable for children.

T-BARDRIN - rectal suppositories for quick and prolonged symptomatic relief in Asthma-Hay Fever.



# WEAK ARCH HERE

Callouses Cramps, Burning, Tenderness



Prescribe Dr. Scholl's Arch Supports in cases requiring mechanical relief from Foot Arch trouble of any kind. The patient will be properly fitted and the Supports adjusted at no extra cost as the condition of the foot improves. This nation-wide service is available at many leading Shoe and Department Stores and at Dr. Scholl's Foot Comfort® Shops in principal cities. For professional literature, write The Scholl Mfg. Co., Inc., 213 West Schiller St., Chicago 10, Ill.

DE Scholl's SUPPORTS



"I just had to talk to you alone, doctor.
and I didn't want to take up your time
at the office."

# Can't Fool Him

To the public, penicillin is a cure-all. A patient at my office had some "local in oil" injected in the sacroiliac region for low back pain. He got off the table with the pain completely gone, and asked. "What did you inject, Doc?"

I told him it was a solution we used for local anesthesia.

"Oh, don't kid me," he retorted, "I'll bet you injected penicillin!"—E.A.Z.

"You know," said the worried patient, "my cousin is in terrible shape. She is pregnant and her doctor tells her she is 4H positive."-L.L.

### High Time, Too

I went to Alabama to begin practice. One of my first patients was a young unmarried girl, obviously pregnant. To determine when the baby might be expected I asked her the date of her last menstruation but was answered with a blank look. I asked her several other questions in an effort to elicit some useful information. To each she replied, "I don't know."

Finally, I said in desperation, "Isn't there anything you can tell me that would help fix a date?"

"Oh yes, doctor," she said brightly, holding her hand about a foot above the floor, "I remember the cotton was about this high."—w.c.



for Balance

ESTROGENS

with

VITAMINS

- A new plus in oral estrogen therapy
- Formula based on the significant interrelationship between vitamin B complex deficiency and the metabolism of estrogens 1, 2
- For the relief of symptoms of estrogen insufficiency associated with subclinical vitamin deficiency
- To restore estrogenic balance and the sense of "fitness" to the menopausal patient

# SESTRAMIN

The Original Formula of Natural Conjugated Estrogens (Equine) with Vitamins B Complex, C and D

Tablets available in two potencies:

SEStramin 10 M (1.25 mg.)

SEStramin 5 M (0.625 mg.)

1 mg.

5 mg.\*

Conjugated estrogens equivalent to oral activity of Sodium Estrone Sulfate 1.25 mg. and 0.625 mg.

In addition both formulas contain:

Brewers' yeast ..... 100 mg.

n

mine hydrochloride . . . 3 mg. (3 MDR)

Riboflavin ...... 2 mg. (1 MDR)

Niscinamide ..... 10 mg.\* d in human nutrition not established.

Ascorbic acid

(Vitamin C) . . . . 25 mg. (% MDR) Vitamin D . . . . . . 500 I. U. (1¼ MDR) MDR-Minimum daily requirements for adults.

Pyridoxine hydrochloride

Calcium pantothenate

Dosage: 1 tablet 1 to 3 times daily, or as directed by the physician. Supplied: Bottles of 20, 100 and 500 tablets.

To be dispensed only by or on the prescription of a physician

Biskind, M. S. and Biskind, G. R.: Endocrinology, 31:109-114 (July) 1942.
 Ashworth, J. and Sutton, D. C.: Arch Int. Med., 69:15-22 (Jan.) 1942

# THE E. L. PATCH COMPANY, Stoneham, Mass.





Before you buy any medical electric instrument. discuss it with your dealer. His advice is invaluable.

## National Electric Instrument Co., Inc. 93-01 Corona Ave., Elmhurst, L.I., N.Y.

"Electric Medical Instruments of Integrity"
ELECTRICATORS, DIAGNOSTIC SETS, NASOPHARYNGOSCOPES, CAUTERY SETS, SOOV CAVITY
SETS, GYOSCOPES, OPHYMALMOSCOPES, MEAD-LIGHTS and other instruments.



### Of Course Not

When I was working for a urologist, we treated an old maid for repeated attacks of cystitis. She was very suspicious and kept wondering where she got the infection. She even imagined that she caught it from her clothing. One day in the course of conversation I asked her if she had a boy friend.

She immediately replied, "Well, there isn't anything like that in this business, I'll have you know."-B.S.

"I feel," remarked the recently delivered OB patient, "as though someone had just moved out."-o.s.

### Sister Mister.

First day of school, while teacher was occupied, the young children were asked to write their most interesting or exciting experience of the summer vacation. All the students wrote long and volubly except Jimmy, who wrote, "Sister Mister.

When all the students had been seated, the teacher, glancing around the room, noticed Jimmy's contribution. "Jimmy," remarked the teacher, "was 'Sister Mister' the most exciting experience you had during the summer?"

"I should say so," said Jimmy. "Only you didn't read it correctly teacher. It says 'Sister Mister Period."-A.S.



"Hey, Nurse! The décolleté is cut too low!"

# 9s There a "GOOD" CIGARETTE?

# THE PHYSICIAN'S VIEWPOINT

gist, ited ious

the she day ked

ess,

was

sk-

or

ca-

ind

Sis-

ed,

m,

y,

er'

ad

nly

If nicotine is contra-indicated, no cigarette qualifies. But if smoking needs to be curtailed only, SANO is the "good" cigarette that paves the way to easy transition.

If the patient finds it difficult and what smoker doesn't?—to reduce the number of cigarettes he or she lights a day, by turning to Sano Cigarettes the nicotine intake is at once reduced to one-half of the amount usually obtained from cigarettes.

A continuing comparative test shows that an average of 51.6 per cent of the nicotine present in the tobacco used in SANO Cigarettes is removed by a special process. The residue of nicotine remaining in the tobacco is less than one per cent.

It is noteworthy that this denicotinizing process does not remove or disturb the essential oils that give tobacco its flavor and aroma. Carefully selected fine tobacco, well aged by maintaining abundant stocks, cured slowly and skillfully blended, assures an enjoyable cigarette that satisfies the expectation of the smoker and meets the demand of the physician for less nicotine for his patient.

SANO Cigarettes are not medicated. They are not a substitute. Pipe tobacco, similarly processed, also is available. It, too, contains less than 1% nicotine.





CARBONATED ALKALINE WATER

Contains those elements necessally to maintain normal alkaline reserve. A palatable, refreshing, zestful drink. Not a laxitive.

KALAK WATER CO. of New York, Inc. 30 Rockefeller Plaza, New York 20, N.Y.





Secretary write for Free Booklet

BABY BATHINETTE CORPORATION ROCHESTER 7, N.Y.



Codeine and hyoscyamus plus ammonium hypophosphite, white pine and tolu in a glycerin base provide sedation of the cough reflex — liquefy mucus. General dosage: Adults 1 to 2 teaspoonfuls every 2 to 3 hrs. Children in proportion. Literature available to physicians.

MARTIN H. SMITH COMPANY

# INDEX TO ADVERTISERS

Ames Company, Inc. Angler Chemical Co. Arlington Chemical Company, The Armour Laboratories, The Baby Bathinette Corporation Barnes, A. C., Company	13 16 12
Angier Chemical Co	16
Arlington Chemical Company, The	120
Armour Laboratories, The	140-14
Baby Bathinette Corporation Barnes, A. C., Company. Bauer & Black. Becton, Dickinson & Co., Bittoher, Corporation, The	
Rarnes A C Company	
Daves & Plack	94 9
Desten Iti-bluson & Co	34-31 37-40
Decton, Dickinson & Co	
Birtcher Corporation, The Borcherdt Malt Extract Company	
Borcherdt Malt Extract Company	110
Breon, George A., & Co	
Brewer & Company, Inc	150
Carand Corporation	
Carnrick, G. W., Company,	
Cereal Institute. Inc.	134-13
Ciba Pharmaceutical Products, Inc	4th Cover
Commercial Colvents Commercian	103
Cutter Laboratories Corporation	9
Cutter Laboratories	14
Devilbiss Company, The	
Dietene Company, The	
Dome Chemicals, Inc	165
Drug Specialties, Inc	
Fleet, C. B., Co., Inc.,,,,	
Fleming-Hall Tobacco Co., Inc.	
Flint, Eaton & Company.	
French R T Co The	
Color Company Inc	
Candalah D B & Co The	100
Birtcher Corporation, The. Borcherdt Mait Extract Company. Breon, George A. & Co. Brewer & Company, Inc. Carand Corporation. Carantek, G. W., Company. Cereal Institute, Inc. Clibs Pharmaceutical Products, Inc. Commercial Solvents Corporation. Cutter Laboratories. Devilibiss Company, The. Dietene Company, The. Dome Chemicals, Inc. Drug Specialites, Inc. Drug Specialites, Inc. Fleet, C. B., Co., Inc. Flening-Hall Tobacco Co., Inc. Flint, Eaton & Company, French, R. T. Co., The. Geodrich, B. F. & Co., The. Hamilton Mfg. Co. Haskell, Charles C., & Co., Inc. Irwin, Neisler & Co. Johnson & Johnson Kalak Water Co. of N. Y. Inc. Knox Gelatine.	
Hashall Charles C	
Haskell, Charles C., & Co., Inc	20
Irwin, Neisler & Co	96-97
Johnson & Johnson	
Kalak Water Co. of N. Y., Inc	168
Johnson & Johnson Kalak Water Co. of N. Y., Inc Knox Gelatine Lakeside Laboratories	
Lakeside Laboratories	
LaMar Laboratories, Inc.	
Leaming Thos & Co Inc	
Tilly Ell & Company	
MacCanger Instrument Company	4
Maket I above topies Inc.	49
McNell Laboratories, Inc	
Mead Johnson & Co	30-31
Lakeside Laboratories LaMar Laboratories, Inc. Leeming, Thos., & Co., Inc. Lilly, Ell & Company.  MacGregor Instrument Company McNeil Laboratories, Inc. Mead Johnson & Co. Merck & Co., Inc. Merrell, The Wm. S. Company. Modern Medicine Annual Musterole.	2nd Cover, 143
Merrell, The Wm. S. Company	2nd Cover, 143
Modern Medicine Annual	156-157
Musterole National Confectioners' Association National Drug Company National Electric Instrument Co., Inc.	160
National Confectioners' Association	
National Drug Company	51
National Electric Instrument Co. Inc.	
Num Specialties	
Num Specialties.  Paravox, Inc.  Parke, Davis & Company.  Patch, The E. L., Company  Pitzer, Chas. S., & Co., Inc.	
Num Specialties  Paravox, Inc.  Parke, Davis & Company.  Patch, The E. L., Company.  Pfizer, Chas. S., & Co., Inc.  Plessner Company, The Paul.	
Num Specialties  Paravox, Inc.  Parke, Davis & Company.  Patch, The E. L., Company.  Pitzer, Chas. S., & Co., Inc.  Plessner Company, The Paul.  Pyramid Rubber Co., The.	
Num Specialties. Paravox, Inc. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S Laboratories.	
Paravoz, no. 2 Company Paravoz, no. 2 Company Patch, The E. J., Company Patch, The E. J., Company Plessner Company, The Paul Pyramid Rubber Co., The Peul P& B Laboratories	
Paravoz, nor de Company Paravoz, nor de Company Patch, The E. J., Company Patch, The E. J., Company Plessner Company, The Paul Pyramid Rubber Co., The Pe & B Laboratories	
Paravoz, nor de Company Paravoz, nor de Company Patch, The E. J., Company Patch, The E. J., Company Plessner Company, The Paul Pyramid Rubber Co., The Pe & B Laboratories	
Paravox, Inc Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company	
Paravox, Inc Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company	
Paravox, Inc Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company	
Paravox, Inc Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company	
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	1622 158 13 165 54 158 46 166 126-127 124 101 24 19,20 151 6-27, 105, 106
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	1622 1158 133 165 144 158 46 1126-127 124 19, 20 16-27, 105, 106 28
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	. 162 . 153 . 133 . 165 . 54 . 158 . 46 . 46 . 166 . 126-127 . 124 . 19 . 20 . 151 . 16-27, 105, 106 . 28 . 15 . 168 . 1
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	162 158 13 165 148 158 46 166 126-127 121 191 101 124 119, 20 16-27, 105, 106 184 184 185 165 167 167 167 167 167 167 167 167 167 167
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	. 162 . 153 . 135 . 145 . 154 . 158 . 46 . 166 . 126-127 . 124 . 19 . 20 . 151 . 148 . 28 . 15 . 161 . 170 . 161
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	162 158 13 165 14 165 158 46 166 166 126 127 19 10 19 20 18 18 18 18 18 18 18 18 18 18 18 18 18
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	162 158 13 165 14 165 158 46 166 166 126 127 19 10 19 20 18 18 18 18 18 18 18 18 18 18 18 18 18
Paravox, Inc Company. Parke, Davis & Company. Patch, The E. L., Company. Pfizer, Chas. S., & Co., Inc. Plessner Company, The Paul. Pyramid Rubber Co., The. P & S. Laboratories. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Reynolds, R. J., Tobacco Company. Revinolds, R. J., Tobacco Company. Robins, A. H., Company, Inc 2	. 162 . 153 . 135 . 145 . 154 . 158 . 46 . 166 . 126-127 . 124 . 19 . 20 . 151 . 148 . 28 . 15 . 161 . 170 . 161
Paravox. Inc	162 153 133 165 143 165 154 158 46 166 126-127 124 191 191 19, 20 188 155 161 170 184 47 44 47
Paravox. Inc	162 158 13 165 14 165 158 166 166 166 126-127 19.10 19.20 16-27, 105, 106 170 164 177 184 48 477 49 164
Paravox. Inc	. 162 . 158 . 13 . 165 . 54 . 158 . 46 . 166 . 126-127 . 121 . 101 . 24 . 101 . 29 . 105 . 106 . 106-27 . 105 . 106 . 107 . 108 . 108 . 170 . 164 . 177 . 44 . 47 . 49 . 164
Paravox. Inc	. 162 . 158 . 13 . 165 . 54 . 158 . 46 . 166 . 126-127 . 121 . 101 . 24 . 101 . 29 . 105 . 106 . 106-27 . 105 . 106 . 107 . 108 . 108 . 170 . 164 . 177 . 44 . 47 . 49 . 164
Paravos, inc.  Paravo	162 158 13 165 148 166 166 166 126-127 191 19 106 17 19 106 18 18 18 18 18 17 19 19 19 19 19 19 19 19 19 19 19 19 19
Paravos, inc.  Paravo	162 153 133 165 144 158 466 166 126-127 124 121 101 101 18, 20 188 155 161 170 164 147 47 49 49 1, 29, 131, 147
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 121 191 19, 20 166-27, 105, 106 177 184 187 187 187 187 187 187 187 187 187 187
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 158 46 168 166 126-127 124 124 19, 20 18 18 170 164 177 44 47 49 49 172 184 172 184 172 184 175 186 186 186 186 186 186 186 186 186 186
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 19 101 19 106-27 105, 106 167 169 167 17 18 18 18 18 18 18 18 18 18 18 18 18 18
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 153 133 165 144 158 466 166 126-127 124 121 101 101 119 18 18 15 161 170 164 147 47 47 49 172 184 185 166 1186 1186 1196 1197 1197 1197 1197 1197 1197 119
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 191 192 16-27, 105, 106 164 177 44 47 49 49 49 49 49 49 49 49 49 49 49 49 49
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 165 13 165 13 165 164 165 166 166 166 166 166 17 17 17 17 17 17 17 17 17 17 17 17 17
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 191 192 16-27, 105, 106 164 177 44 47 49 164 1729, 131, 147 164 168 168 168 168 168 168 168 168 168 168
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 165 13 165 13 165 164 166 166 166 166 17 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 191 192 16-27, 105, 106 16-27, 105, 106 174 184 187 197 194 197 194 197 194 197 194 197 194 197 194 197 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 165 13 165 13 165 166 166 166 126-127 17 19 19 19 19 19 19 19 19 19 19 19 19 19
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 191 192 16-27, 105, 106 16-27, 105, 106 174 184 187 197 194 197 194 197 194 197 194 197 194 197 194 197 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 165 13 165 13 165 166 166 166 126-127 17 19 19 19 19 19 19 19 19 19 19 19 19 19
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 148 166 166 126-127 191 191 192 16-27, 105, 106 16-27, 105, 106 174 184 187 197 194 197 194 197 194 197 194 197 194 197 194 197 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 158 13 165 13 165 158 46 166 166 126-127 19.10 19.20 16-27, 105, 106 17, 106 184 187 197 194 497 194 497 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc. & Company. Paravox. Inc. & Company. Parke, Davis & Company. Patch. The E. L., Company. Pfizer, Chas. S., & Co., Inc. Pflesaner Company. The Paul. Pyramid Rubber Co. The. P & S Laboratories. Raiston-Purina Company. Raiston-Purina Company. Rand Pharmaceuticals Co., Inc. Raymer Pharmacal Company. Raytheon Manufacturing Co. Reed & Carnrick. Raytheon Manufacturing Co. Reed & Carnrick. Royal Metal Mfg. Co. Sanborn Company. Schenley Laboratories, Inc. Schering Corporation. Schenic Jaboratories, Inc. Schering Corporation. Schmid, Jullus, Inc. Scholl Mfg. Co., Inc., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Seamless Rubber Co., The Sex & Kade, Inc. Sharp & Dohme Sherman Laboratories Skiar Smith, Kline & French Laboratories Smith, Martin H., Company. Special Formula Corporation Strauss Laboratories.	162 165 13 165 13 165 166 166 166 166 126-127 101 19, 20 115 168-27, 105, 106 188 28 180 170 184 47 47 47 48 184 177 188 189 180 180 180 180 180 180 180 180 180 180
Paravox. Inc	162 158 13 165 13 165 166 166 166 126-127 101 19, 20 166-27, 105, 106 184 184 187 191 191 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc	162 158 13 165 13 165 166 166 166 126-127 101 19, 20 166-27, 105, 106 184 184 187 191 191 194 195 195 195 195 195 195 195 195 195 195
Paravox. Inc	162 158 13 165 13 165 166 166 166 126-127 101 19, 20 166-27, 105, 106 184 184 187 191 191 194 195 195 195 195 195 195 195 195 195 195
Paravos, Inc. & Company. Paravos, Inc. & Company. Parko, The E. M., Company. Parko, The Paul. Pyennid Rubber Co., The Paul. Pyramid Rubber Co., The Paul. Pyramid Rubber Co., Inc. Rajiston-Purina Company. Randor Pharmacal Company. Raytheon Manufacturing Co. Red & Carnick. Reynolds, R. J., Tobacco Company. Robins, A. H., Company, Inc. 28 Robins, A. H., Company, Inc. 29 Robins, A. H., Company, Inc. 39 Robins, A. H., Company, Inc. 30 Robins, A. M., Company, Inc. 30 Robins, A. Robins, Inc. 31 Robins, A. Robins, Inc. 31 Robins, A. Robins, Inc. 31 Robins, A. Robins, Inc. 32 Robins, A. Robins, Inc. 33 Robins, A. Robins, Inc. 34 Robins, A. Robins,	162 165 133 165 134 165 166 166 166 126-127 161 19, 20 161 181 170 161 184 171 184 184 185 186 186 186 186 186 186 186 186 186 186

# a New Ulmer Specialty Jablet

PANASAL "Ulmer"



PANASAL, the latest of the Ulmer line of fine pharmaceutical tablet specialties, offers a new and effective approach to the treatment of arthritis, rheumatism and rheumatic fever. PANASAL, given orally, permits the high salicylate blood levels formerly obtainable only by intravenous injection under hospital care.

Each PANASAL "Ulmer"
Tablet contains:

Sodium Salicylate ......0.25 Gm. (3.85 gr.)

Para-aminobenzoic Acid
......0.25 Gm. (3.85 gr.)

# INDICATIONS:

Rheumatic fever, arthritic states, rheumatism, neuralgias, myalgias, and such other conditions as are amenable to salicylate therapy. PANASAL "Ulmer" is particularly aimed at assuring high therapeutically effective blood levels of salicylates for the treatment of these conditions.

Write for booklet MM-1149 on The Treatment of Rheumatism, Arthritis and Rheumatic Fever with Panasal "Ulmer." This booklet is available free on request, and gives the rationale of this new treatment method, indications, dosage regimen, etc. A new bibliography is included.

# ULMER PHARMACAL COMPANY

MINNEAPOLIS

Manufacturing Chemists

MINN ESOTA

# Tacts About Conception Control


Use of a diaphragm introducer is favored by many patients who find manual manipulation objectionable or difficult. It facilitates the insertion and correct placement of the diaphragm, as well as its removal.

The "RAMSES" • Diaphragm Introducer provides the following features:

- · Simplicity and convenience in use
- Safety design minimizes possibility of injury to the cervix or accidental insertion into the urethra
- Smooth surface lessens bacterial proliferation makes for easy cleaning
- Ease of removal assured by bluntly hooked end

The "RAMSES" Diaphragm Introducer is supplied in the Physician's Prescription Packet No. 501, without charge







PHYSICIAN'S PRESCRIPTION PACKET NO. 501

A complete unit for conception control. Contains (1) a "RAMSES" Flexible Cushioned Diaphragm of the prescribed size, (2) a "RAMSES" Diaphragm Introducer of corresponding size, and (3) a tube of "RAMSES" Vaginal Jellyt (regular size).

\* The word "RAMSES" is a registered trademark of Julius Schmid, Inc.

† Active Ingredients: Dodecaethyleneglycol Monolaurate 5%; Boric Acid 1%: Alcohol 5%.



Julius Schmid, Snc.

Ty first since 1883

"RAMSES" Vaginal Jelly is accepted by the Council on Pharmacy and Chemistry of the American Medical Association. The "RAMSES" Diaphragm and Diaphragm Introducer are accepted by the Council on Physical Medicine of the American Medical Association.

In the management of arterial hypertension cultivation of sensible habits of living—avoiding unnecessary emotional stress—plays an essential role and aids considerably in the stabilization of pressure on a lower level.

For supplementary medication Theominal, the vasodilator, antispasmodic and sedative, is well suited. Theominal exerts a general tranquilizing effect and thus helps to control temperamental outbursts that may induce dangerous vascular crises.

The average dose is 1 Theominal tablet two or three times daily. With improvement the dose may be reduced or omitted periodically. Each tablet contains 5 grains theobromine and ½ grain Luminal.®

Winthrop-Stearns Inc. New York 13, N. Y. Windsor, Ont.

THEOMINAL for Arterial Hypertension

wominal, trademark reg. U. S. & Conada • Luminal, trademark reg. U. S. & Conada, brand of phenobarbital



# TRASENTINE-PHENOBARBITAL



a powerful antispasmodic...with selective action avoiding undesirable side effects



Effective relief of visceral spasm is generally obtained with Trasentine or Trasentine-Phenobarbital. By its selective action, Trasentine avoids the undesirable side effects of dryness of the mouth and pupillary dilatation frequently produced by belladonna or atropine. These advantages have caused physicians to prescribe more Trasentine and Trasentine-Phenobarbital than probably any other brand of antispasmodic.

• Average adult dose is one or two tablets 3 or 4 times daily as required.

 $T_{\rm RASENTINE-PHENOBARBITAL} - T_{\rm ablets} \ \, ({\rm yellow}) \ \, {\rm contain} \ \, {\rm 5o} \ \, {\rm mg}. \ \, T_{\rm rasentine} \ \, {\rm hydrochloride}$  with 20 mg. phenobarbital, in packages of 100 and 500.

Trasentine – Tablets (white) of 75 mg., in bottles of 100 and 500; also suppositories of 100 mg., and ampuls of 50 mg.

Ciba

PHARMACEUTICAL PRODUCTS, INC., SUMMIT, NEW JERSEY

TRASENTINE (brand of adiphenine)—Trade Mark Reg. U. S. Pat. Off. 2/1431M